



Contents lists available at ScienceDirect

Journal of Quantitative Spectroscopy & Radiative Transfer

journal homepage: www.elsevier.com/locate/jqsrt

Numerical algorithms of the radiative transport equation using rotated reference frames for optical tomography with structured illumination



Manabu Machida

Institute for Medical Photonics Research, Hamamatsu University School of Medicine, Hamamatsu 431-3192, Japan

ARTICLE INFO

Article history:

Received 22 September 2018
 Revised 24 May 2019
 Accepted 11 June 2019
 Available online 12 June 2019

MSC:
 00-01
 99-00

Keywords:

Radiative transport equation
 Optical tomography
 Structured illumination
 Rotated reference frames

ABSTRACT

We consider optical tomography with structured illumination in spatial-frequency domain using the three-dimensional radiative transport equation. Without the diffusion approximation, the radiative transport equation is solved by the technique of rotated reference frames. In addition to the method of rotated reference frames (spherical-harmonic expansion), the three dimensional F_N method is applied to this optical tomography.

© 2019 Elsevier Ltd. All rights reserved.

1. Introduction

Optical tomography is an imaging modality with near-infrared light [1–5]. Compared to the inverse problem of X-ray computed tomography, the inverse problem of optical tomography is more ill-posed since light is strongly scattered. Quite often arrays of optical fibers are used to detect outgoing light on the boundary. One way to improve the resolution of reconstructed images is to increase measured data. Noncontact optical tomography can readily acquire a large number of source-detector pairs [3]. In a typical noncontact optical tomography, a source-detector pair consists of a point source by raster scanning a collimated laser beam and a pixel of a CCD camera [6]. Since the energy density of light in random media such as biological tissue obeys the diffusion equation in the macroscopic regime, in which the propagation distance of light is much larger than the transport mean free path, optical tomography is usually formulated as an inverse problem of the diffusion equation. However, the diffusion approximation breaks in optically thin layers, near boundaries, and in strongly absorbing media. In this mesoscopic regime, in which the propagation distance of light is comparable to the transport mean free path, we need to use the radiative transport equation [7,8], which has angular variables that do not exist in the diffusion equation.

In this paper, we consider noncontact optical tomography based on the radiative transport equation without making the diffusion approximation. We illuminate the boundary (the $x - y$ plane at $z = 0$) of the half space in which the target inhomogeneity is embedded and measure the reflected light on the boundary. The Fourier transform is performed to the data from boundary measurements for source-detector pairs. By the use of spatially modulated beams of structured illumination [9], we can omit the Fourier transform for source positions. Thus the number of measurements can be reduced compared with raster scanned point sources. In this setup of optical tomography with structured illumination, improvement of spatial resolution was observed [10]. Absorbers of different structures can be reconstructed by using bi-dimensional source patterns [11]. The use of angular-dependence of structured light reflectance [12] and dense sampling [13] were proposed. Albeit structured illumination is promising, optical tomography with structured illumination has been mostly limited to the inverse problem of the diffusion equation. Below we will develop transport-based optical tomography for structured illumination. In our optical tomography the absorption coefficient is recovered from boundary data directly measured in the spatial-frequency domain. A reasonably low-cost computation is achieved with algorithms using rotated reference frames.

In the half-space or slab geometry, it is known that plane-wave decomposition is useful [3,14,15]. Different numerical algorithms to compute the Green's function for the radiative transport equation as a sum of plane waves have been developed [16–28]. In

E-mail address: machida@hama-med.ac.jp

We have

$$I_b(\mathbf{r}, \hat{\mathbf{s}}) = \int_{\mathbb{S}_+^2} \int_{\mathbb{R}_+^3} \delta(\boldsymbol{\rho} - \boldsymbol{\rho}') \frac{e^{-|z-z'|}}{\mu'} \delta(\hat{\mathbf{s}} - \hat{\mathbf{s}}') \times \mu' f(\boldsymbol{\rho}', \hat{\mathbf{s}}') \delta(z') d\mathbf{r}' d\hat{\mathbf{s}}' = e^{-z} f(\boldsymbol{\rho}, \hat{\mathbf{s}}), \tag{2.3}$$

for $\hat{\mathbf{s}} \in \mathbb{S}_+^2$, and $I_b = 0$ for $\hat{\mathbf{s}} \in \mathbb{S}_-^2$. Hence

$$S[f](\mathbf{r}, \hat{\mathbf{s}}) = \varpi \int_{\mathbb{S}^2} p(\hat{\mathbf{s}}, \hat{\mathbf{s}}') I_b(\mathbf{r}, \hat{\mathbf{s}}') d\hat{\mathbf{s}}' = \varpi e^{-z} \int_{\mathbb{S}_+^2} p(\hat{\mathbf{s}}, \hat{\mathbf{s}}') f(\boldsymbol{\rho}, \hat{\mathbf{s}}') d\hat{\mathbf{s}}'.$$

Thus the rest is to compute the scattering term given by

$$I_s(\mathbf{r}, \hat{\mathbf{s}}) = \int_{\mathbb{S}^2} \int_{\mathbb{R}_+^3} G(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}', \hat{\mathbf{s}}') S[f](\mathbf{r}', \hat{\mathbf{s}}') d\mathbf{r}' d\hat{\mathbf{s}}'.$$

3. Preliminaries

Both of MRRF (the method of rotated reference frames) and the three-dimensional F_N method are constructed from the three-dimensional Case's method [50]. Polynomials g_l^m and p_l^m are introduced in Section 3.1. Singular eigenfunctions are explained in Section 3.2 and Section 3.3. In Section 3.4, the Green's function is given in terms of singular eigenfunctions.

3.1. Polynomials

Let us begin by introducing h_l ($l = 0, 1, \dots$) as

$$h_l = \begin{cases} 2l + 1 - \varpi \beta_l, & 0 \leq l \leq L, \\ 2l + 1, & l > L. \end{cases}$$

The normalized Chandrasekhar polynomials $g_l^m(\nu)$ ($m \geq 0, l \geq m, \nu \in \mathbb{R}$) are given by the three-term recurrence relation [51,52]

$$\nu h_l g_l^m(\nu) = \sqrt{(l+1)^2 - m^2} g_{l+1}^m(\nu) + \sqrt{l^2 - m^2} g_{l-1}^m(\nu),$$

with the initial term

$$g_m^m(\nu) = \frac{(2m-1)!!}{\sqrt{(2m)!}} = \frac{\sqrt{(2m)!}}{2^m m!}.$$

Moreover we introduce the polynomials $p_l^m(\mu)$ ($m \geq 0, l \geq m$) as

$$p_l^m(\mu) = (-1)^m \sqrt{\frac{(l-m)!}{(l+m)!}} P_l^m(\mu) (1-\mu^2)^{-m/2} = \sqrt{\frac{(l-m)!}{(l+m)!}} \frac{d^m}{d\mu^m} P_l(\mu).$$

The polynomials satisfy the three-term recurrence relation

$$\sqrt{l^2 - m^2} p_{l-1}^m(\mu) - (2l+1)\mu p_l^m(\mu) + \sqrt{(l+1)^2 - m^2} p_{l+1}^m(\mu) = 0.$$

3.2. Singular eigenfunctions for one dimension

In one-dimensional transport theory, singular eigenfunctions $\phi^m(\nu, \mu)$ are given by [53–55]

$$\phi^m(\nu, \mu) = \frac{\varpi \nu}{2} \mathcal{P} \frac{g^m(\nu, \mu)}{\nu - \mu} + \lambda^m(\nu) (1 - \mu^2)^{-|m|} \delta(\nu - \mu),$$

where \mathcal{P} denotes the Cauchy principal value and

$$g^m(\nu, \mu) = \sum_{l=|m|}^L \beta_l p_l^m(\mu) g_l^m(\nu).$$

Here $|m| \leq L$ and $\nu \in \mathbb{R}$ are eigenvalues; ν has discrete values $\pm \nu_j^m$ ($\nu_j^m > 1, j = 0, 1, \dots, M^m - 1$) and the continuous spectrum between -1 and 1 . The number M^m of discrete eigenvalues depends on ϖ and β_l . The function $\lambda^m(\nu)$ is given by

$$\lambda^m(\nu) = 1 - \frac{\varpi \nu}{2} \mathcal{P} \int_{-1}^1 \frac{g^m(\nu, \mu)}{\nu - \mu} (1 - \mu^2)^{|m|} d\mu.$$

Singular eigenfunctions are normalized as

$$\int_{-1}^1 \phi^m(\nu, \mu) (1 - \mu^2)^{|m|} d\mu = 1.$$

We note that

$$g_l^m(\nu) = (-1)^m \sqrt{\frac{(l-m)!}{(l+m)!}} \times \int_{-1}^1 \phi^m(\nu, \mu) (1 - \mu^2)^{|m|/2} P_l^m(\mu) d\mu.$$

Discrete eigenvalues are roots of Λ^m , i.e., $\Lambda^m(\nu_j^m) = 0$, where

$$\Lambda^m(w) = 1 - \frac{\varpi w}{2} \int_{-1}^1 \frac{g^m(w, \mu)}{w - \mu} (1 - \mu^2)^{|m|} d\mu.$$

We have the following orthogonality relations [53–55]

$$\int_{-1}^1 \mu \phi^m(\nu, \mu) \phi^m(\nu', \mu) (1 - \mu^2)^{|m|} d\mu = \mathcal{N}^m(\nu) \delta_{\nu\nu'},$$

where the Kronecker delta $\delta_{\nu\nu'}$ is replaced by the Dirac delta $\delta(\nu - \nu')$ if ν, ν' are in the continuous spectrum. The normalization factor $\mathcal{N}^m(\nu)$ is given by

$$\mathcal{N}^m(\nu) = \begin{cases} \frac{1}{2} (\nu_j^m)^2 g(\nu_j^m, \nu_j^m) \frac{d\Lambda^m(w)}{dw} \Big|_{w=\nu_j^m}, & \nu = \nu_j^m, \\ \nu \Lambda^{m+}(\nu) \Lambda^{m-}(\nu) (1 - \nu^2)^{-|m|}, & \nu \in (-1, 1), \end{cases}$$

where $\Lambda^{m\pm}(\nu) = \lim_{\epsilon \rightarrow 0^+} \Lambda^m(\nu \pm i\epsilon)$.

Finally we introduce

$$\Phi_\nu^m(\hat{\mathbf{s}}) = \phi^m(\nu, \mu) (1 - \mu^2)^{|m|/2} e^{im\varphi}.$$

3.3. Singular eigenfunctions for three dimensions

Let $\psi(\hat{\mathbf{s}}) \in \mathbb{C}$ be a function of $\hat{\mathbf{s}} \in \mathbb{S}^2$. By the operator $\mathcal{R}_{\hat{\mathbf{k}}}$ defined in [21], angles in $\mathcal{R}_{\hat{\mathbf{k}}} \psi(\hat{\mathbf{s}})$ are measured in the rotated reference frame whose z -axis lies in the direction of a unit vector $\hat{\mathbf{k}} \in \mathbb{C}^3$ ($\hat{\mathbf{k}} \cdot \hat{\mathbf{k}} = 1$).

If $\psi(\hat{\mathbf{s}}) \in \mathbb{C}$ has the form

$$\psi(\hat{\mathbf{s}}) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \psi_{lm} Y_{lm}(\hat{\mathbf{s}}), \quad \psi_{lm} \in \mathbb{C},$$

then we have [22,56,57]

$$\begin{aligned} \mathcal{R}_{\hat{\mathbf{k}}} \psi(\hat{\mathbf{s}}) &= \sum_{l=0}^{\infty} \sum_{m=-l}^l \psi_{lm} \\ &\times \sum_{m'=-l}^l e^{-im'\varphi_{\hat{\mathbf{k}}}} d_{m'm}^l(\theta_{\hat{\mathbf{k}}}) Y_{lm'}(\hat{\mathbf{s}}), \\ \mathcal{R}_{\hat{\mathbf{k}}}^{-1} \psi(\hat{\mathbf{s}}) &= \sum_{l=0}^{\infty} \sum_{m=-l}^l \psi_{lm} \\ &\times \sum_{m'=-l}^l e^{im\varphi_{\hat{\mathbf{k}}}} d_{mm'}^l(\theta_{\hat{\mathbf{k}}}) Y_{lm'}(\hat{\mathbf{s}}), \end{aligned}$$

where $\theta_{\hat{\mathbf{k}}}$ and $\varphi_{\hat{\mathbf{k}}}$ are the polar and azimuthal angles of $\hat{\mathbf{k}}$ in the laboratory frame, and $d_{m'm}^l$ are Wigner's d -matrices. In particular, we note that $\mathcal{R}_{\hat{\mathbf{k}}} \hat{\mathbf{s}} \cdot \hat{\mathbf{s}}' = \hat{\mathbf{s}} \cdot \hat{\mathbf{s}}', \mathcal{R}_{\hat{\mathbf{k}}} \mu = \hat{\mathbf{s}} \cdot \hat{\mathbf{k}}$, and

$$\begin{aligned} \mathcal{R}_{\hat{\mathbf{k}}} Y_{lm}^*(\hat{\mathbf{s}}) &= (-1)^m \mathcal{R}_{\hat{\mathbf{k}}} Y_{l,-m}(\hat{\mathbf{s}}) \\ &= (-1)^m \sum_{m'=-l}^l e^{-im'\varphi_{\hat{\mathbf{k}}}} d_{m',-m}^l(\theta_{\hat{\mathbf{k}}}) Y_{lm'}(\hat{\mathbf{s}}) \\ &= \sum_{m'=-l}^l (-1)^{m'} e^{im'\varphi_{\hat{\mathbf{k}}}} d_{m',m}^l(\theta_{\hat{\mathbf{k}}}) Y_{l,-m'}(\hat{\mathbf{s}}) \\ &= \sum_{m'=-l}^l e^{im'\varphi_{\hat{\mathbf{k}}}} d_{m',m}^l(\theta_{\hat{\mathbf{k}}}) Y_{lm'}^*(\hat{\mathbf{s}}). \end{aligned}$$

3.4. The Green's function

For the later purpose of the plane wave decomposition, we give complex unit vectors $\hat{\mathbf{k}}(\nu, \mathbf{q}) \in \mathbb{C}^3$ ($\nu \in \mathbb{R}$, $\mathbf{q} \in \mathbb{R}^2$) by

$$\hat{\mathbf{k}}(\nu, \mathbf{q}) = \begin{pmatrix} -i\nu\mathbf{q} \\ \hat{k}_z(\nu\mathbf{q}) \end{pmatrix},$$

where $q = |\mathbf{q}|$ and

$$\hat{k}_z(\nu\mathbf{q}) = \sqrt{1 + (\nu q)^2}.$$

It turns out that $d_{mm'}^l(\theta_{\hat{\mathbf{k}}(\nu, \mathbf{q})})$ are functions of νq . We write

$$d_{mm'}^l[i\tau(\nu\mathbf{q})] = d_{mm'}^l(\theta_{\hat{\mathbf{k}}(\nu, \mathbf{q})}).$$

These d -matrices are computed using recurrence relations [21,44,50]. To calculate $d_{mm'}^l[i\tau(\nu\mathbf{q})]$, we take square roots such that $0 \leq \arg(\sqrt{z}) < \pi$ for all $z \in \mathbb{C}$ [23,44]. We have

$$\varphi_{\hat{\mathbf{k}}(\nu, \mathbf{q})} = \begin{cases} \varphi_{\mathbf{q}} + \pi, & \text{for } \nu > 0, \\ \varphi_{\mathbf{q}}, & \text{for } \nu < 0, \end{cases}$$

where $\varphi_{\mathbf{q}}$ is the polar angle of \mathbf{q} .

Let us consider the following homogeneous equation.

$$(\hat{\mathbf{s}} \cdot \nabla + 1)I(\mathbf{r}, \hat{\mathbf{s}}) = \varpi \int_{\mathbb{S}^2} p(\hat{\mathbf{s}}, \hat{\mathbf{s}}') I(\mathbf{r}, \hat{\mathbf{s}}') d\hat{\mathbf{s}}'.$$

Let us consider the Fourier transform of the specific intensity:

$$\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) = \int_{\mathbb{R}^2} e^{-i\mathbf{q} \cdot \boldsymbol{\rho}} I(\mathbf{r}, \hat{\mathbf{s}}) d\boldsymbol{\rho}.$$

Then the Fourier transform $\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}})$ satisfies

$$\begin{aligned} (\mu\partial_z + i\boldsymbol{\omega} \cdot \mathbf{q} + 1)\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) &= \left(1 - \frac{\hat{\mathbf{s}} \cdot \hat{\mathbf{k}}}{\nu}\right)\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) \\ &= \varpi \int_{\mathbb{S}^2} p(\hat{\mathbf{s}}, \hat{\mathbf{s}}') \tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}') d\hat{\mathbf{s}}', \end{aligned} \quad (3.1)$$

where $\boldsymbol{\omega} \in \mathbb{R}^2$ was defined such that $\hat{\mathbf{s}} = (\boldsymbol{\omega}, \mu)$. The solutions are given by

$$\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) = e^{-\hat{k}_z(\nu\mathbf{q})z/\nu} \mathcal{R}_{\hat{\mathbf{k}}(\nu, \mathbf{q})} \Phi_{\nu}^m(\hat{\mathbf{s}}).$$

Three-dimensional singular eigenfunctions are obtained as

$$\begin{aligned} \mathcal{R}_{\hat{\mathbf{k}}} \phi^m(\nu, \mu) &= \frac{\varpi \nu}{2} \mathcal{P} \frac{g^m(\nu, \hat{\mathbf{s}} \cdot \hat{\mathbf{k}})}{\nu - \hat{\mathbf{s}} \cdot \hat{\mathbf{k}}} \\ &\quad + \lambda^m(\nu) (1 - \nu^2)^{-|m|} \delta(\nu - \hat{\mathbf{s}} \cdot \hat{\mathbf{k}}), \end{aligned}$$

where $\hat{\mathbf{k}} = \hat{\mathbf{k}}(\nu, \mathbf{q})$.

The following orthogonality relation holds.

$$\begin{aligned} \int_{\mathbb{S}^2} \mu (\mathcal{R}_{\hat{\mathbf{k}}(\nu, \mathbf{q})} \Phi_{\nu}^m(\hat{\mathbf{s}})) (\mathcal{R}_{\hat{\mathbf{k}}(\nu', \mathbf{q})} \Phi_{\nu'}^{m'*}(\hat{\mathbf{s}})) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\nu\mathbf{q}) \mathcal{N}(\nu) \delta_{\nu\nu'} \delta_{mm'}. \end{aligned}$$

Since the general solution is given by the sum of a particular solution and a linear combination of eigenmodes, we can write the Green's function as [7]

$$\begin{aligned} G(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}_0, \hat{\mathbf{s}}_0) &= G_{\text{free}}(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}_0, \hat{\mathbf{s}}_0) \\ &\quad + \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q} \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}_0)} \\ &\quad \times \sum_{m=-L}^L \left[\sum_{j=0}^{M^m-1} \hat{A}^m(\nu_j^m) \mathcal{R}_{\hat{\mathbf{k}}(\nu_j^m, \mathbf{q})} \Phi_j^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu_j^m\mathbf{q})z/\nu_j^m} \right. \\ &\quad \left. + \int_0^1 \hat{A}^m(\nu) \mathcal{R}_{\hat{\mathbf{k}}(\nu, \mathbf{q})} \Phi_{\nu}^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu\mathbf{q})z/\nu} d\nu \right] d\mathbf{q}, \end{aligned}$$

where $G_{\text{free}}(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}_0, \hat{\mathbf{s}}_0)$ is the free-space Green's function and $\hat{A}^m(\nu_j^m, \mathbf{q})$, $\hat{A}^m(\nu, \mathbf{q})$ are some coefficients which are determined from the boundary conditions. We note that the free-space Green's function or the fundamental solution is obtained as [50]

$$\begin{aligned} G_{\text{free}}(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}_0, \hat{\mathbf{s}}_0) \\ = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q} \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}_0)} \tilde{G}_{\text{free}}(z, \hat{\mathbf{s}}; z_0, \hat{\mathbf{s}}_0; \mathbf{q}) d\mathbf{q}, \end{aligned}$$

where

$$\begin{aligned} \tilde{G}_{\text{free}}(z, \hat{\mathbf{s}}; z_0, \hat{\mathbf{s}}_0; \mathbf{q}) &= \sum_{m=-L}^L \left[\sum_{j=0}^{M^m-1} \frac{1}{2\pi \hat{k}_z(\nu_j^m\mathbf{q}) \mathcal{N}(\nu_j^m)} \right. \\ &\quad \times \mathcal{R}_{\hat{\mathbf{k}}(\nu_j^m, \mathbf{q})} \Phi_{j\pm}^m(\hat{\mathbf{s}}) \Phi_{j\pm}^{m*}(\hat{\mathbf{s}}_0) e^{-\hat{k}_z(\nu_j^m\mathbf{q})|z-z_0|/\nu_j^m} \\ &\quad \left. + \int_0^1 \frac{1}{2\pi \hat{k}_z(\nu\mathbf{q}) \mathcal{N}(\nu)} \mathcal{R}_{\hat{\mathbf{k}}(\pm\nu, \mathbf{q})} \Phi_{\pm\nu}^m(\hat{\mathbf{s}}) \Phi_{\pm\nu}^{m*}(\hat{\mathbf{s}}_0) \right. \\ &\quad \left. \times e^{-\hat{k}_z(\nu\mathbf{q})|z-z_0|/\nu} d\nu \right]. \end{aligned}$$

Upper signs are chosen for $z > z_0$ and lower signs are chosen for $z < z_0$. The Fourier transform of $G(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}_0, \hat{\mathbf{s}}_0)$ is defined similarly.

Let us consider the Fourier transform \tilde{I}_s of I_s , which is given by

$$\tilde{I}_s(\mathbf{r}, \hat{\mathbf{s}}) = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q} \cdot \boldsymbol{\rho}} \tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) d\mathbf{q}.$$

We can express $\tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}})$ in terms of the Green's function and singular eigenfunctions as

$$\begin{aligned} \tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) &= \int_0^\infty \int_{\mathbb{S}^2} \tilde{G}(z, \hat{\mathbf{s}}; z', \hat{\mathbf{s}}'; \mathbf{q}) S[\tilde{f}](\mathbf{q}, z', \hat{\mathbf{s}}') d\hat{\mathbf{s}}' dz' \\ &= \int_0^\infty \int_{\mathbb{S}^2} \tilde{G}_{\text{free}}(z, \hat{\mathbf{s}}; z', \hat{\mathbf{s}}'; \mathbf{q}) S[\tilde{f}](\mathbf{q}, z', \hat{\mathbf{s}}') d\hat{\mathbf{s}}' dz' \\ &\quad + \sum_{m=-L}^L \left[\sum_{j=0}^{M^m-1} A^m(\nu_j^m) \mathcal{R}_{\hat{\mathbf{k}}(\nu_j^m, \mathbf{q})} \Phi_j^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu_j^m\mathbf{q})z/\nu_j^m} \right. \\ &\quad \left. + \int_0^1 A^m(\nu) \mathcal{R}_{\hat{\mathbf{k}}(\nu, \mathbf{q})} \Phi_{\nu}^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu\mathbf{q})z/\nu} d\nu \right], \end{aligned} \quad (3.2)$$

with some coefficients $A^m(\nu_j^m, \mathbf{q})$, $A^m(\nu, \mathbf{q})$. We note that $\tilde{S}[f] = S[\tilde{f}]$, where $\tilde{S}(\mathbf{q}, z, \hat{\mathbf{s}}) = \int_{\mathbb{R}^2} e^{-i\mathbf{q} \cdot \boldsymbol{\rho}} S(\mathbf{r}, \hat{\mathbf{s}}) d\boldsymbol{\rho}$ and $\tilde{f}(\mathbf{q}, \hat{\mathbf{s}}) = \int_{\mathbb{R}^2} e^{-i\mathbf{q} \cdot \boldsymbol{\rho}} f(\boldsymbol{\rho}, \hat{\mathbf{s}}) d\boldsymbol{\rho}$.

4. Method of rotated reference frames

The method of rotated reference frames was first proposed by Markel [22]. Here, we formulate the method making the relation to Case's method clear.

In the method of rotated reference frames, we expand the singular eigenfunction as

$$\Phi_{\nu}^m(\hat{\mathbf{s}}) = \sum_{l=|m|}^N C_l^m(\nu) Y_{lm}(\hat{\mathbf{s}}).$$

$$\begin{aligned} \tilde{f}^{(1)}(\mathbf{q}, \hat{\mathbf{s}}) &= 1, \\ \tilde{f}^{(2)}(\mathbf{q}, \hat{\mathbf{s}}) &= \delta(\hat{\mathbf{s}} - \hat{\mathbf{z}}) = \sum_{l=0}^N \sqrt{\frac{2l+1}{4\pi}} Y_{l0}(\hat{\mathbf{s}}), \end{aligned} \quad (4.3)$$

where we used $Y_{lm}(\hat{\mathbf{z}}) = \sqrt{(2l+1)/(4\pi)}\delta_{m0}$. Correspondingly we write $F_{M\nu}^{(+)}(\mathbf{q}) = F_{M\nu}^{(+i)}(\mathbf{q})$ ($i = 1, 2$). By multiplying $Y_{lm}^*(\hat{\mathbf{s}})$ on both sides and integrating in S_+^2 , we have

$$\begin{aligned} \sum_{M=-L}^L \sum_{\nu>0} \sum_{l'=|m|}^N \sqrt{\frac{2l'+1}{h_{l'}}} \mathcal{B}_{l'}^m \langle l' | \psi_\nu(M) \rangle d_{mM}^{l'} [i\tau(\nu q)] \\ \times \begin{vmatrix} F_{M\nu}^{(+1)}(\mathbf{q}) \\ F_{M\nu}^{(+2)}(\mathbf{q}) \end{vmatrix} = \delta_{m0} \begin{vmatrix} \sqrt{4\pi} \mathcal{B}_{l_0}^0 \\ \sum_{l'=0}^N \sqrt{\frac{2l'+1}{4\pi}} \mathcal{B}_{l'}^0 \end{vmatrix}. \end{aligned}$$

Here we defined

$$\begin{aligned} \mathcal{B}_{l'}^m &= \int_{S_+^2} Y_{lm}^*(\hat{\mathbf{s}}) Y_{l'm}(\hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &= \frac{1}{2} \sqrt{\frac{(2l+1)(2l'+1)(l-m)!(l'-m)!}{(l+m)!(l'+m)!}} \\ &\quad \times \int_0^1 P_l^m(\mu) P_{l'}^m(\mu) d\mu, \end{aligned}$$

and used

$$\begin{aligned} \int_0^1 P_l(\mu) d\mu &= \frac{2}{\sqrt{2l+1}} \mathcal{B}_{l_0}^0 \\ &= \begin{cases} 1, & l=0, \\ 0, & \text{even } l (\neq 0), \\ (-1)^{(l-1)/2} \frac{l!!}{l(l+1)(l-1)!!}, & \text{odd } l. \end{cases} \end{aligned}$$

The fact that $(-1)^M F_{-M,\nu}^{(+)}$ satisfies the same equation as $F_{M\nu}^{(+)}$ implies

$$(-1)^M F_{-M,\nu}^{(+)} = F_{M\nu}^{(+)},$$

where we note that [22] $\langle l | \psi_{-\nu}(M) \rangle = (-1)^l \langle l | \psi_\nu(M) \rangle$. We obtain

$$\begin{aligned} \sum_{M=0}^L \sum_{\nu>0} \sum_{l'=|m|}^N \sqrt{\frac{2l'+1}{h_{l'}}} \mathcal{B}_{l'}^m \langle l' | \psi_\nu(M) \rangle \\ \times [d_{mM}^{l'} [i\tau(\nu q)] + (1 - \delta_{M0})(-1)^M d_{m,-M}^{l'} [i\tau(\nu q)]] \\ \times \begin{vmatrix} F_{M\nu}^{(+1)}(\mathbf{q}) \\ F_{M\nu}^{(+2)}(\mathbf{q}) \end{vmatrix} = \delta_{m0} \begin{vmatrix} \sqrt{4\pi} \mathcal{B}_{l_0}^0 \\ \sum_{l'=0}^N \sqrt{\frac{2l'+1}{4\pi}} \mathcal{B}_{l'}^0 \end{vmatrix}. \end{aligned}$$

Thus $F_{M\nu}^{(+i)}(\mathbf{q})$ ($0 \leq M \leq N$) are obtained from the linear system,

$$\mathcal{M}(\mathbf{q}) \mathbf{F}^{(+i)} = \mathbf{v}^{(+i)}, \quad i = 1, 2,$$

where the matrix $\mathcal{M}(\mathbf{q})$ is defined as

$$\begin{aligned} \{\mathcal{M}(\mathbf{q})\}_{lm,M\nu} &= \sum_{l'=|m|}^N \sqrt{\frac{2l'+1}{h_{l'}}} \mathcal{B}_{l'}^m \langle l' | \psi_\nu(M) \rangle \\ &\quad \times [d_{mM}^{l'} [i\tau(\nu q)] \\ &\quad + (1 - \delta_{M0})(-1)^M d_{m,-M}^{l'} [i\tau(\nu q)]], \end{aligned}$$

and vectors $\mathbf{F}^{(+i)}, \mathbf{v}^{(+i)}$ are given by

$$\{\mathbf{F}^{(+i)}\}_{M\nu} = F_{M\nu}^{(+i)}, \quad i = 1, 2,$$

$$\{\mathbf{v}^{(+1)}\}_{lm} = \delta_{m0} \sqrt{4\pi} \mathcal{B}_{l_0}^0,$$

$$\{\mathbf{v}^{(+2)}\}_{lm} = \delta_{m0} \sum_{l'=0}^N \sqrt{\frac{2l'+1}{4\pi}} \mathcal{B}_{l'}^0.$$

The matrix $\mathcal{M}(\mathbf{q})$ becomes square when $L = M, l = |m| + 2\alpha - 1, \alpha = 1, \dots, \lfloor (N - |m| + 1)/2 \rfloor$ (see Remark 5.1 below). That is,

$$\begin{aligned} I^{(0)}(\mathbf{r}, \hat{\mathbf{s}}) &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q}\cdot\rho} \sum_{M=-L}^L \sum_{\nu>0} e^{-\hat{k}_z(\nu q)z/\nu} F_{M\nu}^{(+i)}(\mathbf{q}) \\ &\quad \times \sum_{l=0}^N \sum_{m=-l}^l \sqrt{\frac{2l+1}{h_l}} \langle l | \psi_\nu(M) \rangle d_{mM}^l [i\tau(\nu q)] \\ &\quad \times (-1)^m e^{-im\varphi_q} Y_{lm}(\hat{\mathbf{s}}) d\mathbf{q}, \quad i = 1, 2. \end{aligned}$$

Now, we note that the ballistic term (2.3) is calculated as

$$\begin{aligned} I_b(\mathbf{r}, \hat{\mathbf{s}}) &= e^{-z} \begin{vmatrix} f^{(1)}(\rho, \hat{\mathbf{s}}) \\ f^{(2)}(\rho, \hat{\mathbf{s}}) \end{vmatrix} \\ &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q}\cdot\rho} e^{-z} \begin{vmatrix} \sqrt{4\pi} Y_{00}(\hat{\mathbf{s}}) \\ \sum_{l=0}^N \sqrt{\frac{2l+1}{4\pi}} Y_{l0}(\hat{\mathbf{s}}) \end{vmatrix} d\mathbf{q}. \end{aligned}$$

Therefore,

$$\begin{aligned} I_s(\mathbf{r}, \hat{\mathbf{s}}) &= I^{(0)}(\mathbf{r}, \hat{\mathbf{s}}) - I_b(\mathbf{r}, \hat{\mathbf{s}}) \\ &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q}\cdot\rho} \sum_{m=-N}^N \sum_{l=|m|}^N \begin{vmatrix} \gamma_{lm}^{(1)}(\mathbf{q}, z) \\ \gamma_{lm}^{(2)}(\mathbf{q}, z) \end{vmatrix} Y_{lm}(\hat{\mathbf{s}}) d\mathbf{q}. \end{aligned}$$

Here we defined

$$\begin{aligned} \gamma_{lm}^{(1)}(\mathbf{q}, z) &= \sqrt{\frac{2l+1}{h_l}} (-1)^m e^{-im\varphi_q} \hat{\gamma}_{lm}^{(1)}(\mathbf{q}, z) \\ &\quad - \sqrt{4\pi} \delta_{l0} \delta_{m0} e^{-z}, \\ \gamma_{lm}^{(2)}(\mathbf{q}, z) &= \sqrt{\frac{2l+1}{h_l}} (-1)^m e^{-im\varphi_q} \hat{\gamma}_{lm}^{(2)}(\mathbf{q}, z) \\ &\quad - \delta_{m0} \sqrt{\frac{2l+1}{4\pi}} e^{-z}, \end{aligned}$$

where

$$\begin{aligned} \hat{\gamma}_{lm}^{(i)}(\mathbf{q}, z) &= \sum_{M=-L}^L \sum_{\nu>0} F_{M\nu}^{(+i)}(\mathbf{q}) e^{-\hat{k}_z(\nu q)z/\nu} \langle l | \psi_\nu(M) \rangle \\ &\quad \times d_{mM}^l [i\tau(\nu q)], \quad i = 1, 2. \end{aligned}$$

5. Three-dimensional F_N method

By combining the F_N method and the technique of rotated reference frames, we can establish a numerical scheme for the three-dimensional radiative transport equation with anisotropic scattering.

Let us introduce the notation

$$\sum'_{lm} = \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor}.$$

For $\hat{\mathbf{s}} \in S_+^2$, we expand \tilde{I}_s as

$$\tilde{I}_s(\mathbf{q}, 0, -\hat{\mathbf{s}}) = \sum'_{lm} c_{lm}(\mathbf{q}) Y_{lm}(\hat{\mathbf{s}}),$$

$$\tilde{I}_s(\mathbf{q}, z, -\hat{\mathbf{s}}) = \sum'_{lm} b_{lm}(\mathbf{q}, z) Y_{lm}(\hat{\mathbf{s}}),$$

$$\tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) = \sum'_{lm} a_{lm}(\mathbf{q}, z) Y_{lm}(\hat{\mathbf{s}}).$$

We define ξ_j^m ($-L \leq m \leq L$) as

$$\xi_j^m = \begin{cases} v_j^m, & j = 0, 1, \dots, M^m - 1, \\ \cos\left(\frac{\pi}{2} \frac{j - M^m + 1}{N_{\text{row}}^m - M^m + 1}\right), & j = M^m, \dots, N_{\text{row}}^m - 1, \end{cases} \quad (5.1)$$

where

$$N_{\text{row}}^m = \left\lfloor \frac{N - |m|}{2} \right\rfloor + 1. \tag{5.2}$$

We drop the superscript m if there is no confusion.

Remark 5.1. How to discretize ν in the continuous spectrum depends on N, L . We selected ξ_j^m in (5.1) because we set $N = L$ in the numerical calculation below.

Furthermore we introduce

$$\begin{aligned} \mathcal{I}_{lm}^m(\xi_j, \mathbf{q}) &= \int_{\mathbb{S}^2} \left(\mathcal{R}_{\hat{\mathbf{k}}(\xi_j, \mathbf{q})} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) Y_{lm}(\hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &= \sqrt{(2l+1)\pi} (-1)^{m'} e^{im\varphi_k} d_{mm'}^l [i\tau(\xi_j q)] g_l^{m'}(\xi_j), \end{aligned}$$

where $\mathcal{I}_{lm}^m(\xi_j, \mathbf{q}) = 0$ for $|m'| > l$. We have

$$\begin{aligned} \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, \mathbf{0}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = L_1^{m'}[\tilde{f}](-\xi_j, \mathbf{q}, \mathbf{0}), \end{aligned} \tag{5.3}$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, \mathbf{0}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi_j q) \mathcal{N}^{m'}(\xi_j) A^{m'}(\xi_j), \end{aligned} \tag{5.4}$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = L_1^{m'}[\tilde{f}](-\xi_j, \mathbf{q}, z), \end{aligned} \tag{5.5}$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi_j q) \mathcal{N}^{m'}(\xi_j) A^{m'}(\xi_j) e^{-\hat{k}_z(\xi_j q)z/\xi_j} \\ + L_2^{m'}[\tilde{f}](\xi_j, \mathbf{q}, z), \end{aligned} \tag{5.6}$$

where

$$\begin{aligned} L_1^{m'}[\tilde{f}](-\xi_j, \mathbf{q}, z) &= \int_z^\infty \int_{\mathbb{S}^2} \left(\mathcal{R}_{\hat{\mathbf{k}}(-\xi_j, \mathbf{q})} \Phi_{-\xi_j}^{m'*}(\hat{\mathbf{s}}') \right) \\ &\quad \times e^{-\hat{k}_z(\xi_j q)(z-z')/\xi_j} \tilde{S}[\tilde{f}](\mathbf{q}, z', \hat{\mathbf{s}}') d\hat{\mathbf{s}}' dz' \\ &= e^{-z} \frac{\varpi \xi_j}{\xi_j + \hat{k}_z(\xi_j q)} \sum_{l=0}^L \sum_{m=-l}^l \frac{\beta_l}{2l+1} \\ &\quad \times \mathcal{I}_{lm}^m(-\xi_j, \mathbf{q}) \int_{\mathbb{S}_+^2} \tilde{f}(\mathbf{q}, \hat{\mathbf{s}}) Y_{lm}^*(\hat{\mathbf{s}}) d\hat{\mathbf{s}}, \end{aligned}$$

$$\begin{aligned} L_2^{m'}[\tilde{f}](\xi_j, \mathbf{q}, z) &= \int_0^z \int_{\mathbb{S}^2} \left(\mathcal{R}_{\hat{\mathbf{k}}(\xi_j, \mathbf{q})} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}') \right) \\ &\quad \times e^{-\hat{k}_z(\xi_j q)(z-z')/\xi_j} \tilde{S}[\tilde{f}](\mathbf{q}, z', \hat{\mathbf{s}}') d\hat{\mathbf{s}}' dz' \\ &= \left(e^{-\hat{k}_z(\xi_j q)z/\xi_j} - e^{-z} \right) \frac{\varpi \xi_j}{\xi_j - \hat{k}_z(\xi_j q)} \sum_{l=0}^L \sum_{m=-l}^l \frac{\beta_l}{2l+1} \\ &\quad \times \mathcal{I}_{lm}^m(\xi_j, \mathbf{q}) \int_{\mathbb{S}_+^2} \tilde{f}(\mathbf{q}, \hat{\mathbf{s}}) Y_{lm}^*(\hat{\mathbf{s}}) d\hat{\mathbf{s}}. \end{aligned}$$

In particular for $\tilde{f}^{(i)}(\mathbf{q}, \hat{\mathbf{s}})$ ($i = 1, 2$) in (4.3), we have

$$\begin{aligned} L_1^{m'}[\tilde{f}](-\xi_j, \mathbf{q}, z) &= e^{-z} \frac{\varpi \xi_j}{2(\xi_j + \hat{k}_z(\xi_j q))} \sum_{l=|m'|}^L (-1)^l \beta_l \\ &\quad \times d_{0m'}^l [i\tau(\xi_j q)] g_l^{m'}(\xi_j) \begin{cases} \frac{4\pi}{\sqrt{2l+1}} \mathcal{B}_{l0}^0, & i = 1, \\ 1, & i = 2, \end{cases} \end{aligned}$$

$$\begin{aligned} L_2^{m'}[\tilde{f}](\xi_j, \mathbf{q}, z) &= \left(e^{-\hat{k}_z(\xi_j q)z/\xi_j} - e^{-z} \right) \frac{\varpi \xi_j}{2(\xi_j - \hat{k}_z(\xi_j q))} \\ &\quad \times \sum_{l=|m'|}^L \beta_l d_{0m'}^l [i\tau(\xi_j q)] g_l^{m'}(\xi_j) \\ &\quad \times \begin{cases} \frac{4\pi}{\sqrt{2l+1}} \mathcal{B}_{l0}^0, & i = 1, \\ 1, & i = 2. \end{cases} \end{aligned}$$

By noticing $g_l^{-m}(\nu) = (-1)^m g_l^m(\nu)$, we see that $L_1^{m'}[\tilde{f}]$ and $L_2^{m'}[\tilde{f}]$ are independent of $\varphi_{\mathbf{q}}$ and independent of the sign of m' . From (5.3) we obtain

$$\begin{aligned} -(-1)^{m'} \int_{\mathbb{S}_+^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, \mathbf{0}, -\hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = L_1^{m'}[\tilde{f}](-\xi_j, \mathbf{q}, \mathbf{0}), \end{aligned}$$

where we used $\Phi_{-\xi_j}^{m'}(\hat{\mathbf{s}}) = (-1)^{m'} \Phi_{\xi_j}^{m'}(-\hat{\mathbf{s}})$. We introduce

$$\begin{aligned} \mathcal{J}_{lm}^{(\pm)jm'} &= e^{-im\varphi_{\mathbf{q}}} \int_{\mathbb{S}_\pm^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}(-\xi_j, \mathbf{q})} \Phi_{-\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) Y_{lm}(\hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &= (-1)^{l+1} e^{-im\varphi_{\mathbf{q}}} \int_{\mathbb{S}_\mp^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}(-\xi_j, \mathbf{q})} \Phi_{-\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) Y_{lm}(\hat{\mathbf{s}}) d\hat{\mathbf{s}}. \end{aligned}$$

Explicit expressions of $\mathcal{J}_{lm}^{(\pm)jm'}$ are found as follows [21].

$$\begin{aligned} \mathcal{J}_{lm}^{(+)jm'} &= \hat{k}_z(\xi_j q) \sqrt{\frac{\pi}{2l+1}} (-1)^m d_{mm'}^l [i\tau(\xi_j q)] \\ &\quad \times \left(\sqrt{(l+1)^2 - m'^2} g_{l+1}^{m'}(\xi_j) + \sqrt{l^2 - m'^2} g_{l-1}^{m'}(\xi_j) \right) \\ &\quad - i \frac{|\xi_j q|}{2} \sqrt{\frac{\pi}{2l+1}} (-1)^m \sum_{m''=-l}^l d_{mm''}^l [i\tau(\xi_j q)] \\ &\quad \times \left[\delta_{m'', m'-1} \left(\sqrt{(l-m'')(l-m')} g_{l-1}^{m''}(\xi_j) \right. \right. \\ &\quad \left. \left. - \sqrt{(l+m'+1)(l+m')} g_{l+1}^{m''}(\xi_j) \right) \right. \\ &\quad \left. + \delta_{m'', m'+1} \left(\sqrt{(l-m'+1)(l-m')} g_{l+1}^{m''}(\xi_j) \right. \right. \\ &\quad \left. \left. - \sqrt{(l+m'')(l+m')} g_{l-1}^{m''}(\xi_j) \right) \right] \\ &\quad - \mathcal{J}_{lm}^{(-)jm'}, \end{aligned}$$

where we used $g_l^m(-\nu) = (-1)^{l+m} g_l^m(\nu)$, and

$$\begin{aligned} \mathcal{J}_{lm}^{(-)jm'} &= \frac{\varpi \xi_j}{2} (-1)^{l+1} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} [\text{sgn}(m')]^{m'} \\ &\quad \times \frac{\sqrt{(2|m'|)!}}{(2|m'| - 1)!!} \sum_{m''=-|m'|}^{|m'|} (-1)^{m''} \sqrt{\frac{(|m'| - m'')!}{(|m'| + m'')!}} \\ &\quad \times d_{m'' - m'}^{|m'|} [i\tau(\xi_j q)] \\ &\quad \times \int_{\mathbb{S}_+^2} \frac{g^{m'} \left(-\xi_j, \hat{k}_z(\xi_j q) \mu + i\xi_j q \sqrt{1 - \mu^2} \cos \varphi \right)}{\xi_j + \hat{k}_z(\xi_j q) \mu + i\xi_j q \sqrt{1 - \mu^2} \cos \varphi} \\ &\quad \times \mu P_{|m'|}^{m''}(\mu) P_l^m(\mu) e^{i(m+m'')\varphi} d\hat{\mathbf{s}}. \end{aligned}$$

When numerically evaluating the above integral over $\hat{\mathbf{s}}$, we use the Gauss-Legendre quadrature for μ and trapezoidal rule for φ . We note that

$$\mathcal{J}_{l, -m}^{(\pm)j, -m'} = (-1)^m \mathcal{J}_{lm}^{(\pm)jm'}. \tag{5.7}$$

For each ξ_j, \mathbf{q} , we have

$$\sum'_{lm} \mathcal{J}_{lm}^{(+jm')} c_{lm}(\mathbf{q}) e^{im\varphi_q} = -L_1^{m'} [\tilde{f}](-\xi_j, \mathbf{q}, 0).$$

Using (5.7) and the fact that $L_1^{-m'} = L_1^{m'}$, we have

$$\begin{aligned} & \sum'_{lm} \mathcal{J}_{lm}^{(+jm')} c_{lm}(\mathbf{q}) e^{im\varphi_q} \\ &= \sum'_{lm} \mathcal{J}_{lm}^{(+j,-m')} c_{lm}(\mathbf{q}) e^{im\varphi_q} \\ &= \sum'_{lm} \mathcal{J}_{l,-m}^{(+j,-m')} c_{l,-m}(\mathbf{q}) e^{-im\varphi_q} \\ &= \sum'_{lm} \mathcal{J}_{lm}^{(+jm')} c_{l,-m}(\mathbf{q}) (-1)^m e^{-im\varphi_q}. \end{aligned}$$

The above relation implies

$$c_{lm}(\mathbf{q}) = \hat{c}_{lm}(q) e^{-im\varphi_q}, \quad \hat{c}_{l,-m}(q) = (-1)^m \hat{c}_{lm}(q).$$

That is,

$$\begin{aligned} & \sum_{m=0}^N \sum_{l=|m|, |m|+2, \dots} \left[\mathcal{J}_{lm}^{(+jm')} + (1 - \delta_{m0}) (-1)^m \mathcal{J}_{l,-m}^{(+jm')} \right] \\ & \times \hat{c}_{lm}(q) = -L_1^{m'} [\tilde{f}](-\xi_j, \mathbf{q}, 0). \end{aligned}$$

We obtain from (5.4) and (5.6)

$$\begin{aligned} & \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &= e^{-\hat{k}_z(\xi_j q)z/\xi_j} \int_{\mathbb{S}^2} \mu \left(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi_j}^{m'*}(\hat{\mathbf{s}}) \right) \tilde{I}_s(\mathbf{q}, 0, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ & \quad + L_2^{m'} [\tilde{f}](\xi_j, \mathbf{q}, z). \end{aligned} \tag{5.8}$$

Together with (5.5), we have

$$\begin{aligned} & \sum'_{lm} (-1)^{l+1} \mathcal{J}_{lm}^{(-jm')} a_{lm}(\mathbf{q}, z) e^{im\varphi_q} \\ &= \sum'_{lm} \mathcal{J}_{lm}^{(+jm')} b_{lm}(\mathbf{q}, z) e^{im\varphi_q} + L_1^{m'} [\tilde{f}](-\xi_j, \mathbf{q}, z), \\ & \sum'_{lm} (-1)^{m'} \mathcal{J}_{lm}^{(+jm')} a_{lm}(\mathbf{q}, z) e^{im\varphi_q} \\ & \quad - \sum'_{lm} (-1)^{l+m'+1} \mathcal{J}_{lm}^{(-jm')} b_{lm}(\mathbf{q}, z) e^{im\varphi_q} \\ &= e^{-\hat{k}_z(\xi_j q)z/\xi_j} (-1)^{m'} \left(\sum'_{lm} (-1)^l \mathcal{J}_{lm}^{(-jm')} c_{lm}(\mathbf{q}) e^{im\varphi_q} \right) \\ & \quad + L_2^{m'} [\tilde{f}](\xi_j, \mathbf{q}, z), \end{aligned}$$

where we used $Y_{lm}(-\hat{\mathbf{s}}) = (-1)^l Y_{lm}(\hat{\mathbf{s}})$. Let us define

$$\begin{aligned} \nu_1^{jm'} &= -L_1^{m'} [\tilde{f}](-\xi_j, \mathbf{q}, z), \\ \nu_2^{jm'} &= e^{-\hat{k}_z(\xi_j q)z/\xi_j} \left(\sum'_{lm} (-1)^l \mathcal{J}_{lm}^{(-jm')} c_{lm}(\mathbf{q}) \right) \\ & \quad + (-1)^{m'} L_2^{m'} [\tilde{f}](\xi_j, \mathbf{q}, z). \end{aligned}$$

We can show that $\nu_1^{jm'}$ and $\nu_2^{jm'}$ are independent of φ_q and satisfy $\nu_1^{j,-m'} = \nu_1^{jm'}$ and $\nu_2^{j,-m'} = \nu_2^{jm'}$. Similarly to $c_{lm}(\mathbf{q})$ we have

$$\begin{aligned} a_{lm}(\mathbf{q}, z) &= \hat{a}_{lm}(q, z) e^{-im\varphi_q}, \\ \hat{a}_{l,-m}(q, z) &= (-1)^m \hat{a}_{lm}(q, z), \\ b_{lm}(\mathbf{q}, z) &= \hat{b}_{lm}(q, z) e^{-im\varphi_q}, \\ \hat{b}_{l,-m}(q, z) &= (-1)^m \hat{b}_{lm}(q, z). \end{aligned}$$

We write

$$\mathcal{M}_{jm',lm}^{(11)} = \mathcal{M}_{jm',lm}^{(22)}$$

$$\begin{aligned} &= (-1)^l \mathcal{J}_{lm}^{(-jm')} + (1 - \delta_{m0}) (-1)^{l+m} \mathcal{J}_{l,-m}^{(-jm')}, \\ & \mathcal{M}_{jm',lm}^{(12)} = \mathcal{M}_{jm',lm}^{(21)} \\ &= \mathcal{J}_{lm}^{(+jm')} + (1 - \delta_{m0}) (-1)^m \mathcal{J}_{l,-m}^{(+jm')}. \end{aligned}$$

We obtain

$$\left(\begin{array}{c|c} \mathcal{M}_{jm',lm}^{(11)} & \mathcal{M}_{jm',lm}^{(12)} \\ \hline \mathcal{M}_{jm',lm}^{(21)} & \mathcal{M}_{jm',lm}^{(22)} \end{array} \right) \begin{pmatrix} \hat{a}_{lm}(q, z) \\ \hat{b}_{lm}(q, z) \end{pmatrix} = \begin{pmatrix} \nu_1^{jm'} \\ \nu_2^{jm'} \end{pmatrix}.$$

By setting $N=L$ and choosing $0 \leq m' \leq L$, $0 \leq j \leq N_{\text{row}}^{m'} - 1$, $0 \leq m \leq N$, $l = m + 2\alpha$, $0 \leq \alpha \leq \lfloor \frac{N-m}{2} \rfloor$, the above matrix \mathcal{M} becomes a square matrix. Otherwise the above linear system can be solved by singular value decomposition.

In Appendix A, the three-dimensional F_N method for the slab geometry is explained.

6. Optical tomography with structured illumination

We here explain the set up of our optical tomography. The data function introduced in Section 6.2 is obtained for spatially modulated incident beams in Section 6.1, and $\eta(\mathbf{r})$ is reconstructed according to the inversion formula (6.9) in Section 6.3.

6.1. Structured illumination

Let us consider structured illumination in the half space, i.e., the incoming beam is given by

$$\begin{aligned} f(\boldsymbol{\rho}, \hat{\mathbf{s}}) &= F_{B_0}(\boldsymbol{\rho}, \hat{\mathbf{s}}) \\ &= I_0 [1 + A_0 \cos(\mathbf{q}_0 \cdot \boldsymbol{\rho} + B_0)] \delta(\hat{\mathbf{s}} - \hat{\mathbf{s}}_0), \end{aligned}$$

where $\mu_0 \in (0, 1]$ is the cosine of the polar angle of $\hat{\mathbf{s}}_0$, I_0 is the amplitude, A_0 is the modulation depth, and B_0 is the phase of the source. Since

$$\begin{aligned} & \frac{2F_0(\boldsymbol{\rho}, \hat{\mathbf{s}}) - (1 - i\sqrt{3})F_{-2\pi/3}(\boldsymbol{\rho}, \hat{\mathbf{s}}) - (1 + i\sqrt{3})F_{2\pi/3}(\boldsymbol{\rho}, \hat{\mathbf{s}})}{3A_0} \\ &= f_{q_0}(\boldsymbol{\rho}, \hat{\mathbf{s}}), \end{aligned}$$

where

$$f_{q_0}(\boldsymbol{\rho}, \hat{\mathbf{s}}) = I_0 e^{iq_0 \cdot \boldsymbol{\rho}} \delta(\hat{\mathbf{s}} - \hat{\mathbf{s}}_0), \quad \mu_0 \in (0, 1],$$

we will use $f(\boldsymbol{\rho}, \hat{\mathbf{s}}) = f_{q_0}(\boldsymbol{\rho}, \hat{\mathbf{s}})$ for the boundary condition [48]. Then we have

$$\begin{aligned} \tilde{f}(\mathbf{q}, \hat{\mathbf{s}}) &= \int_{\mathbb{R}^2} e^{-iq \cdot \boldsymbol{\rho}} f(\boldsymbol{\rho}, \hat{\mathbf{s}}) d\boldsymbol{\rho} \\ &= (2\pi)^2 I_0 \delta(\mathbf{q} - \mathbf{q}_0) \delta(\hat{\mathbf{s}} - \hat{\mathbf{s}}_0). \end{aligned} \tag{6.1}$$

We measure the exitance or hemispheric flux J_+ defined as follows on the boundary.

$$J_+(\boldsymbol{\rho}) = \int_{\mathbb{S}_+^2} \mu I(\boldsymbol{\rho}, 0, -\hat{\mathbf{s}}) d\hat{\mathbf{s}}.$$

Moreover we consider the Fourier transform of η :

$$\tilde{\eta}(\mathbf{q}, z) = \int_{\mathbb{R}^2} e^{-iq \cdot \boldsymbol{\rho}} \eta(\boldsymbol{\rho}, z) d\boldsymbol{\rho}.$$

Since the reconstructed $\tilde{\eta}$ is regularized, the reconstructed η is given by

$$\begin{aligned} \eta(\boldsymbol{\rho}, z) &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{iq \cdot \boldsymbol{\rho}} \chi_{\Omega_B}(\mathbf{q}) \tilde{\eta}(\mathbf{q}, z) d\mathbf{q} \\ &=: \eta_B(\boldsymbol{\rho}, z), \end{aligned} \tag{6.2}$$

where Ω_B is a subdomain in the first Brillouin zone (see Section 6.2) and $\chi_{\Omega_B}(\mathbf{q})$ is the characteristic function such that $\chi_{\Omega_B}(\mathbf{q}) = 1$ for $\mathbf{q} \in \Omega_B$ and $\chi_{\Omega_B}(\mathbf{q}) = 0$ otherwise.

6.2. The data function

We refer to $D(\boldsymbol{\rho})$ below as the data function.

$$\begin{aligned} D(\boldsymbol{\rho}) &= J_+^{(0)}(\boldsymbol{\rho}) - J_+(\boldsymbol{\rho}) \\ &= \int_{\mathbb{S}_+^2} \mu [I^{(0)}(\boldsymbol{\rho}, 0, -\hat{\mathbf{s}}) - I(\boldsymbol{\rho}, 0, -\hat{\mathbf{s}})] d\hat{\mathbf{s}}. \end{aligned}$$

Within the first Born approximation, the data function $D(\boldsymbol{\rho})$ is given by

$$\begin{aligned} D(\boldsymbol{\rho}) &= \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} \left[\int_{\mathbb{S}_+^2} \mu G(\boldsymbol{\rho}, 0, -\hat{\mathbf{s}}; \mathbf{r}', \hat{\mathbf{s}}') d\hat{\mathbf{s}} \right] \\ &\quad \times \eta_B(\mathbf{r}') I^{(0)}(\mathbf{r}', \hat{\mathbf{s}}') d\mathbf{r}' d\hat{\mathbf{s}}'. \end{aligned} \quad (6.3)$$

We note that $D(\boldsymbol{\rho})$ is obtained through the exitance measured on the boundary. We can reconstruct η_B by solving the linear inverse problem in (6.3).

Suppose that there are $(2N_d + 1) \times (2N_d + 1)$ detectors on grid points $\boldsymbol{\rho} = (x_i, y_j)$ ($i, j = -N_d, \dots, N_d$) with spacing h_d . We consider the Fourier transform

$$\tilde{D}(\mathbf{q}) = \sum_{\boldsymbol{\rho}} e^{-i\mathbf{q} \cdot \boldsymbol{\rho}} D(\boldsymbol{\rho}).$$

Noting the Poisson summation formula

$$\sum_{\boldsymbol{\rho}} e^{i\mathbf{q} \cdot \boldsymbol{\rho}} = \left(\frac{2\pi}{h_d} \right)^2 \sum_{\mathbf{p}} \delta(\mathbf{q} + \mathbf{p}),$$

where \mathbf{p} denotes reciprocal lattice points $(2\pi i/h_d, 2\pi j/h_d)$ ($i, j \in \mathbb{Z}$). We note that Ω_B is in the first Brillouin zone $[-\pi/h_d, \pi/h_d] \times [-\pi/h_d, \pi/h_d]$. If $\mathbf{q} - \mathbf{q}_0 \in \Omega_B$, we obtain

$$\begin{aligned} \tilde{D}(\mathbf{q}) &= \frac{\mu_0 I_0}{h_d^2} \int_0^\infty \int_{\mathbb{S}^2} \left[\int_{\mathbb{S}_+^2} \mu \tilde{G}^*(z', -\hat{\mathbf{s}}'; 0, \hat{\mathbf{s}}_0; \mathbf{q}) d\hat{\mathbf{s}} \right] \\ &\quad \times \tilde{\eta}_B(\mathbf{q} - \mathbf{q}_0, z') \tilde{G}(z', \hat{\mathbf{s}}'; 0, \hat{\mathbf{s}}_0; \mathbf{q}_0) d\hat{\mathbf{s}}' dz'. \end{aligned} \quad (6.4)$$

To compute the right-hand side of (6.4), we recall $f^{(1)}, f^{(2)}$ in (4.2) and introduce

$$I^{(i)}(\mathbf{r}, \hat{\mathbf{s}}) = \int_{\mathbb{S}_+^2} \int_{\mathbb{R}^3} G(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}', \hat{\mathbf{s}}') \mu' f^{(i)}(\boldsymbol{\rho}', \hat{\mathbf{s}}') \delta(z') d\mathbf{r}' d\hat{\mathbf{s}}',$$

for $i = 1, 2$ and its Fourier transform

$$\tilde{I}^{(i)}(\mathbf{q}, z, \hat{\mathbf{s}}) = \int_{\mathbb{S}_+^2} \tilde{G}(z, \hat{\mathbf{s}}; 0, \hat{\mathbf{s}}'; \mathbf{q}) \mu' \tilde{f}^{(i)}(\mathbf{q}, \hat{\mathbf{s}}') d\hat{\mathbf{s}}',$$

for $i = 1, 2$. We note that $\tilde{f}^{(i)}$ are given in (4.3) and $I^{(i)}(\mathbf{r}, \hat{\mathbf{s}})$ satisfies (2.1) with the boundary source $f^{(i)}(\boldsymbol{\rho}, \hat{\mathbf{s}})$ ($i = 1, 2$). Thus we can write $\tilde{D}(\mathbf{q})$ as

$$\begin{aligned} \tilde{D}(\mathbf{q}) &= \frac{I_0}{h_d^2} \int_0^\infty \int_{\mathbb{S}^2} \tilde{I}^{(1)*}(\mathbf{q}, z', -\hat{\mathbf{s}}') \tilde{\eta}_B(\mathbf{q} - \mathbf{q}_0, z') \\ &\quad \times \tilde{I}^{(2)}(\mathbf{q}_0, z', \hat{\mathbf{s}}') d\hat{\mathbf{s}}' dz'. \end{aligned} \quad (6.5)$$

By redefining \mathbf{q} as $\mathbf{q} - \mathbf{q}_0 \rightarrow \mathbf{q}$ in (6.5), we have

$$\mathcal{D}(\mathbf{q}_0, \mathbf{q}) = \int_0^\infty K(\mathbf{q}_0, z; \mathbf{q}) \tilde{\eta}_B(\mathbf{q}, z) dz, \quad \mathbf{q} \in \Omega_B, \quad (6.6)$$

where

$$\mathcal{D}(\mathbf{q}_0, \mathbf{q}) = \tilde{D}(\mathbf{q} + \mathbf{q}_0) \frac{h_d^2}{I_0},$$

and

$$K(\mathbf{q}_0, z; \mathbf{q}) = \int_{\mathbb{S}^2} \tilde{I}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z, -\hat{\mathbf{s}}) \tilde{I}^{(2)}(\mathbf{q}_0, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}}.$$

We obtain

$$K(\mathbf{q}_0, z; \mathbf{q}) = e^{-z} \int_{\mathbb{S}_+^2} \tilde{I}_s^{(1)*}(\mathbf{q} + \mathbf{q}_0, z, -\hat{\mathbf{s}}) \tilde{f}^{(2)}(\mathbf{q}_0, \hat{\mathbf{s}}) d\hat{\mathbf{s}}$$

$$\begin{aligned} &+ e^{-z} \int_{\mathbb{S}_+^2} \tilde{I}_s^{(2)}(\mathbf{q}_0, z, -\hat{\mathbf{s}}) \tilde{f}^{(1)*}(\mathbf{q} + \mathbf{q}_0, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &+ \int_{\mathbb{S}^2} \tilde{I}_s^{(1)*}(\mathbf{q} + \mathbf{q}_0, z, -\hat{\mathbf{s}}) \tilde{I}_s^{(2)}(\mathbf{q}_0, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}}. \end{aligned} \quad (6.7)$$

Thus the quality of reconstruction of η in this inverse problem is determined by the kernel K in (6.6). So far in most research including [48], K has been computed within the diffusion approximation. In the present paper, we will calculate K directly from the radiative transport equation.

Below, we will numerically compute $\tilde{I}_s^{(1)}, \tilde{I}_s^{(2)}$ using MRRF and the three-dimensional F_N method.

6.3. Reconstruction by MRRF

We expand $\tilde{I}_s^{(1)}$ and $\tilde{I}_s^{(2)}$ with spherical harmonics:

$$\begin{aligned} \tilde{I}_s^{(1)}(\mathbf{q}, z', -\hat{\mathbf{s}}') &= \sum_{m=-N}^N \sum_{l=|m|}^N \gamma_{lm}^{(1)}(\mathbf{q}, z') Y_{lm}(-\hat{\mathbf{s}}'), \quad \hat{\mathbf{s}}' \in \mathbb{S}^2, \\ \tilde{I}_s^{(2)}(\mathbf{q}_0, z', \hat{\mathbf{s}}') &= \sum_{m=-N}^N \sum_{l=|m|}^N \gamma_{lm}^{(2)}(\mathbf{q}_0, z') Y_{lm}(\hat{\mathbf{s}}'), \quad \hat{\mathbf{s}}' \in \mathbb{S}^2. \end{aligned}$$

Using MRRF, we obtain

$$\begin{aligned} K(\mathbf{q}_0, z; \mathbf{q}) &= e^{-z} \sum_{l=0}^N \sum_{l'=0}^N (-1)^l \gamma_{l0}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z) \mathcal{B}_{ll'}^0 \\ &\quad + e^{-z} \sqrt{4\pi} \sum_{l=0}^N (-1)^l \gamma_{l0}^{(2)}(\mathbf{q}_0, z) \mathcal{B}_{l0}^0 \\ &\quad + \sum_{m=-N}^N \sum_{l=|m|}^N (-1)^l \gamma_{lm}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z) \gamma_{lm}^{(2)}(\mathbf{q}_0, z), \end{aligned} \quad (6.8)$$

where we used $Y_{lm}(-\hat{\mathbf{s}}) = (-1)^l Y_{lm}(\hat{\mathbf{s}})$ and $\int_{\mathbb{S}_+^2} Y_{lm}(\hat{\mathbf{s}}) d\hat{\mathbf{s}} = \sqrt{4\pi} \delta_{m0} \mathcal{B}_{l0}^0$. Hence,

$$\begin{aligned} K(\mathbf{q}_0, z; \mathbf{q}) &= e^{-z} \sum_{l=0}^N \sum_{l'=0}^N (-1)^l \\ &\quad \times \left[\sqrt{\frac{2l+1}{h_l}} \hat{\gamma}_{l0}^{(1)*}(|\mathbf{q} + \mathbf{q}_0|, z) - e^{-z} \sqrt{4\pi} \delta_{l0} \right] \mathcal{B}_{ll'}^0 \\ &\quad + e^{-z} \sqrt{4\pi} \sum_{l=0}^N (-1)^l \left[\sqrt{\frac{2l+1}{h_l}} \hat{\gamma}_{l0}^{(2)}(q_0, z) - e^{-z} \sqrt{\frac{2l+1}{4\pi}} \right] \mathcal{B}_{l0}^0 \\ &\quad + \sum_{l=0}^N (-1)^l \left[\sqrt{\frac{2l+1}{h_l}} \hat{\gamma}_{l0}^{(1)*}(|\mathbf{q} + \mathbf{q}_0|, z) - e^{-z} \sqrt{4\pi} \delta_{l0} \right] \\ &\quad \times \left[\sqrt{\frac{2l+1}{h_l}} \hat{\gamma}_{l0}^{(2)}(q_0, z) - e^{-z} \sqrt{\frac{2l+1}{4\pi}} \right] \\ &\quad + 2 \sum_{m=1}^N \sum_{l=m}^N (-1)^l \frac{2l+1}{h_l} \hat{\gamma}_{lm}^{(1)*}(|\mathbf{q} + \mathbf{q}_0|, z) \hat{\gamma}_{lm}^{(2)}(q_0, z) \\ &\quad \times \cos(m(\varphi_{\mathbf{q} + \mathbf{q}_0} - \varphi_{\mathbf{q}_0})). \end{aligned}$$

Viewing (6.6) as a linear matrix-vector equation we can express (6.6) as

$$|\mathcal{D}(\mathbf{q})\rangle = \hat{K}(\mathbf{q}) |\tilde{\eta}_B(\mathbf{q})\rangle,$$




Fig. 1. Reconstruction by MRRF. Three panels show the reconstruction of $\delta\mu_a(\mathbf{r})/\max_{\mathbf{r}\in\mathbb{R}^2}(\delta\mu_a(\mathbf{r}))$ in planes parallel to the $x-y$ plane at depths, from the left, 1 cm, 2 cm, and 3 cm. The point absorber is placed at the center of the plane on the z -axis at the depth 2 cm. The field of view is $5.1\text{ cm} \times 5.1\text{ cm}$.

where $\langle \mathbf{q}_0 | \mathcal{D}(\mathbf{q}) \rangle = \mathcal{D}(\mathbf{q}_0, \mathbf{q})$, $\langle \mathbf{q}_0 | \hat{K}(\mathbf{q}) | z \rangle = K(\mathbf{q}_0, z; \mathbf{q})$, and $\langle z | \tilde{\eta}_B(\mathbf{q}) \rangle = \tilde{\eta}_B(\mathbf{q}, z)$. We can compute $|\tilde{\eta}_B(\mathbf{q})\rangle$ with singular value decomposition. We obtain

$$\tilde{\eta}_B(\mathbf{q}, z) = \langle z | \hat{K}^+(\mathbf{q}) | \mathcal{D}(\mathbf{q}) \rangle,$$

where \hat{K}^+ is the pseudoinverse such that

$$\hat{K}^+ = \hat{K}^\dagger (\hat{K} \hat{K}^\dagger)^{-1}_{\text{reg}}.$$

Here \dagger denotes the Hermitian conjugate and reg means that the pseudoinverse is regularized as is explained below. Let $\sigma_j^2(\mathbf{q})$ and $|v_j(\mathbf{q})\rangle$ be the eigenvalues and eigenvectors of the matrix $\hat{M}(\mathbf{q})$ whose $\mathbf{q}_0 - \mathbf{q}'_0$ element is given by

$$\langle \mathbf{q}_0 | \hat{M}(\mathbf{q}) | \mathbf{q}'_0 \rangle = \int_0^\infty K(\mathbf{q}_0, z; \mathbf{q}) K^*(\mathbf{q}'_0, z; \mathbf{q}) dz.$$

If we use the truncated SVD and take only singular values greater than a threshold value σ_0 as regularization, η is reconstructed as

$$\eta(\mathbf{r}) = \frac{1}{(2\pi)^2} \int_{\Omega_B} e^{i\mathbf{q}\cdot\mathbf{r}} \sum_{\substack{j \\ \sigma_j > \sigma_0}} \sigma_j(\mathbf{q})^{-2} \sum_{\mathbf{q}_0} \langle v_j(\mathbf{q}) | \mathbf{q}_0 \rangle \times \mathcal{D}(\mathbf{q}_0, \mathbf{q}) \sum_{\mathbf{q}'_0} K^*(\mathbf{q}'_0, z; \mathbf{q}) \langle \mathbf{q}'_0 | v_j(\mathbf{q}) \rangle d\mathbf{q}. \quad (6.9)$$

The resulting reconstruction is shown in Fig. 1.

6.4. Reconstruction by the three-dimensional F_N method

We expand $\tilde{\eta}_S^{(1)}$ and $\tilde{\eta}_S^{(2)}$ with spherical harmonics:

$$\begin{aligned} & \tilde{\eta}_S^{(1)}(\mathbf{q}, z', -\hat{\mathbf{s}}') \\ &= \begin{cases} \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} b_{lm}^{(1)}(\mathbf{q}, z') Y_{lm}(\hat{\mathbf{s}}'), & \hat{\mathbf{s}}' \in \mathbb{S}_+^2, \\ \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} a_{lm}^{(1)}(\mathbf{q}, z') Y_{lm}(-\hat{\mathbf{s}}'), & \hat{\mathbf{s}}' \in \mathbb{S}_-^2, \end{cases} \end{aligned}$$

and

$$\begin{aligned} & \tilde{\eta}_S^{(2)}(\mathbf{q}_0, z', \hat{\mathbf{s}}') \\ &= \begin{cases} \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} a_{lm}^{(2)}(\mathbf{q}_0, z') Y_{lm}(\hat{\mathbf{s}}'), & \hat{\mathbf{s}}' \in \mathbb{S}_+^2, \\ \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} b_{lm}^{(2)}(\mathbf{q}_0, z') Y_{lm}(-\hat{\mathbf{s}}'), & \hat{\mathbf{s}}' \in \mathbb{S}_-^2. \end{cases} \end{aligned}$$

Note that $Y_{lm}(-\hat{\mathbf{s}}') = (-1)^l Y_{lm}(\hat{\mathbf{s}}')$. Since $\hat{\mathbf{s}}_0 \in \mathbb{S}_+^2$ and $-\hat{\mathbf{s}} \in \mathbb{S}_-^2$ in (6.4), the ballistic terms do not contribute to $\tilde{D}(\mathbf{q})$. The kernel in (6.7) is obtained as

$$\begin{aligned} & K(\mathbf{q}_0, z; \mathbf{q}) \\ &= e^{-z} \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} b_{lm}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z) Y_{lm}^*(\hat{\mathbf{z}}) \end{aligned}$$

$$\begin{aligned} & + e^{-z} \sqrt{4\pi} \sum_{\substack{\alpha=0 \\ l=2\alpha}}^{\lfloor N/2 \rfloor} b_{l0}^{(2)}(\mathbf{q}_0, z) \mathcal{B}_{l0}^0 \\ & + \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} \sum_{\substack{\alpha'=0 \\ l'=|m|+2\alpha'}}^{\lfloor (N-|m|)/2 \rfloor} \left[a_{lm}^{(2)}(\mathbf{q}_0, z) b_{l'm}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z) \right. \\ & \left. + b_{lm}^{(2)}(\mathbf{q}_0, z) a_{l'm}^{(1)*}(\mathbf{q} + \mathbf{q}_0, z) \right] \mathcal{B}_{ll'}^m. \quad (6.10) \end{aligned}$$

Thus the kernel K is obtained using the three-dimensional F_N method. We can rewrite (6.10) as

$$\begin{aligned} & K(\mathbf{q}_0, z; \mathbf{q}) = e^{-z} \sum_{\substack{\alpha=0 \\ l=2\alpha}}^{\lfloor N/2 \rfloor} \sqrt{\frac{2l+1}{4\pi}} \hat{b}_{l0}^{(2)*}(|\mathbf{q} + \mathbf{q}_0|, z) \\ & + e^{-z} \sqrt{4\pi} \sum_{\substack{\alpha=0 \\ l=2\alpha}}^{\lfloor N/2 \rfloor} \hat{b}_{l0}^{(2)}(q_0, z) \mathcal{B}_{l0}^0 \\ & + \sum_{m=0}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-m)/2 \rfloor} \sum_{\substack{\alpha'=0 \\ l'=|m|+2\alpha'}}^{\lfloor (N-m)/2 \rfloor} \left[\hat{a}_{lm}^{(2)}(q_0, z) \hat{b}_{l'm}^{(1)*}(|\mathbf{q} + \mathbf{q}_0|, z) \right. \\ & \left. + \hat{b}_{lm}^{(2)}(q_0, z) \hat{a}_{l'm}^{(1)*}(|\mathbf{q} + \mathbf{q}_0|, z) \right] \\ & \times \left[\delta_{m0} + 2(1 - \delta_{m0}) \cos(m(\varphi_{\mathbf{q}+\mathbf{q}_0} - \varphi_{\mathbf{q}_0})) \right] \mathcal{B}_{ll'}^m, \end{aligned}$$

where we used $(-1)^m Y_{l-m}^*(\hat{\mathbf{s}}) = Y_{lm}(\hat{\mathbf{s}})$. The absorption inhomogeneity $\eta(\mathbf{r})$ is obtained from (6.9).

7. Simulation

To show that the numerical schemes developed in Section 4 and Section 5 are capable of optical tomography, we perform optical tomography with structured illumination. We consider a random medium which has the following optical properties.

$$\mu_a = 0.05\text{ cm}^{-1}, \quad \mu_s = 100\text{ cm}^{-1}, \quad g = 0.9005. \quad (7.1)$$

For these optical parameters, the transport mean free path is $\ell^* = (\mu_t - \mu_s g)^{-1} = 1\text{ mm}$, which is typical in biological tissue.

For the formulation in Section 6, the forward data $\mathcal{D}(\mathbf{q}_0, \mathbf{q})$ is calculated by the diffusion equation, which is an approximation of the radiative transport equation. In this way, we can avoid inverse crime. We assume N_a point targets at positions $\mathbf{r}_a^{(i)} = (\rho_a^{(i)}, z_a^{(i)})$ ($i = 1, \dots, N_a$). Appendix B is devoted to the computation of the data function $\mathcal{D}(\mathbf{q}_0, \mathbf{q})$.

We place detectors on a square lattice of spacing h_d . Detector positions are specified by

$$(x_i, y_j) = (ih_d, jh_d), \quad i, j = -N_d, \dots, N_d.$$

We set

$$h_d = 1\text{ mm}, \quad N_d = 25.$$

Let $q^{(x)(i)}$, $q^{(y)(j)}$ denote the x - and y -components of \mathbf{q} . We put $W = [2(N_d + N_s) + 1]h_d$, and

$$q^{(x)(i)} = \frac{2\pi}{W} i, \quad q^{(y)(j)} = \frac{2\pi}{W} j, \quad i, j = -N_d, \dots, N_d.$$

For \mathbf{q}_0 we use

$$q_0^{(x)(i)} = \frac{2\pi}{W} i, \quad q_0^{(y)(j)} = \frac{2\pi}{W} j, \quad i, j = -N_s, \dots, N_s.$$

We set

$$N_s = 10.$$

Note that $q_0^{(x)(i+1)} - q_0^{(x)(i)} \approx 0.1\text{ mm}^{-1}$. In this demonstration we chose $L = N = 9$, $l_0 = 1$, and $\hat{\mathbf{s}}_0 = \hat{\mathbf{z}}$. Reconstruction is done by the formula (6.9). In both cases of MRRF and the three-dimensional F_N method, about 10 singular values were used for the truncated SVD.




Fig. 2. Reconstruction by the three-dimensional F_N method. Three panels show the reconstruction of $\delta\mu_a(\mathbf{r})/\max_{\mathbf{r}\in\mathbb{R}_+^3}(\delta\mu_a(\mathbf{r}))$ in planes parallel to the $x-y$ plane at depths, from the left, 1 cm, 2 cm, and 3 cm. The point absorber is placed at the center of the plane on the z -axis at the depth 2 cm. The field of view is $5.1\text{ cm} \times 5.1\text{ cm}$.




Fig. 3. Reconstruction by MRRF. Four panels show the reconstruction of $\delta\mu_a(\mathbf{r})/\max_{\mathbf{r}\in\mathbb{R}_+^3}(\delta\mu_a(\mathbf{r}))$ in planes parallel to the $x-y$ plane at depths, from the left, 5 mm, 10 mm, 15 mm, and 20 mm. The point absorbers are placed at $(10\text{ mm}, 10\text{ mm}, 10\text{ mm})$ and $(-10\text{ mm}, -10\text{ mm}, 15\text{ mm})$. The field of view is $5.1\text{ cm} \times 5.1\text{ cm}$.

7.1. A point absorber

First we put $N_a = 1$. Let us assume that a point absorber is embedded at $x_a^{(1)} = y_a^{(1)} = 0$, $z_a^{(1)} = 20\text{ mm}$.

Fig. 1 shows the reconstruction by MRRF. In Fig. 1, $\delta\mu_a(\mathbf{r})/\max_{\mathbf{r}\in\mathbb{R}_+^3}(\delta\mu_a(\mathbf{r}))$ is plotted in planes parallel to the $x-y$ plane at different depths z . In the middle panel for $z = 2\text{ cm}$, the absorber placed 2 cm away from the surface is reconstructed while the reconstructed $\eta \approx 0$ in the left panel for $z = 1\text{ cm}$ and the right panel for $z = 3\text{ cm}$.

Fig. 2 shows the reconstruction by the three-dimensional F_N method. The results are almost identical, and the target is successfully reconstructed.

7.2. Two point absorbers

Next we consider two absorbers ($N_a = 2$): One at $\mathbf{r}_a^{(1)} = (10\text{ mm}, 10\text{ mm}, 10\text{ mm})$ and the other at $\mathbf{r}_a^{(2)} = (-10\text{ mm}, -10\text{ mm}, 15\text{ mm})$. Reconstructed images are shown in Figs. 3 and 4 for MRRF and the three-dimensional F_N method, respectively. Tomographic images show that reconstruction of multiple targets is more difficult than that of a single target. In Fig. 3 for MRRF, the target at $\mathbf{r}_a^{(2)}$ is almost invisible whereas the target at $\mathbf{r}_a^{(1)}$ is almost invisible in Fig. 4 for the three-dimensional F_N method.

To investigate the robustness of reconstruction, we added 3% Gaussian noise to the data function $\mathcal{D}(\mathbf{q}_0, \mathbf{q})$. The results are shown in Figs. 5 and 6.

8. Concluding remarks

Since the aim of this paper is to present novel numerical algorithms of the radiative transport equation for optical tomography in spatial-frequency domain, we mainly focused on how to




Fig. 4. Reconstruction by the three-dimensional F_N method. Four panels show the reconstruction of $\delta\mu_a(\mathbf{r})/\max_{\mathbf{r}\in\mathbb{R}_+^3}(\delta\mu_a(\mathbf{r}))$ in planes parallel to the $x-y$ plane at depths, from the left, 5 mm, 10 mm, 15 mm, and 20 mm. The point absorbers are placed at $(10\text{ mm}, 10\text{ mm}, 10\text{ mm})$ and $(-10\text{ mm}, -10\text{ mm}, 15\text{ mm})$. The field of view is $5.1\text{ cm} \times 5.1\text{ cm}$.




Fig. 5. Same as Fig. 3 but 3% Gaussian noise is added.




Fig. 6. Same as Fig. 4 but 3% Gaussian noise is added.

compute $\tilde{r}^{(1)}, \tilde{r}^{(2)}$ in (6.5) and reconstructions are done for simple point targets. Reconstructions of targets of more complicated shapes will be necessary in the future study. In particular, it is a future problem to show the superiority of the transport-based optical tomography over the diffusion-based optical tomography in spatial-frequency domain. Since structured illumination is quite often considered in shallow regions where the diffusion approximation breaks [49], the radiative transport equation is expected to be important for structured illumination.

It is straightforward to extend the three-dimensional F_N method for the half-space to the slab as is described in Appendix A. In [44], MRRF was formulated in a slab. Thus our optical tomography can be similarly formulated in the slab geometry.

Although the first Born approximation is employed in this paper (see (2.2)), recently optical tomography with higher-order nonlinear terms has been developed by the use of a recursion algorithm for the inverse Born series of the radiative transport equation [58]. The present schemes for calculating the kernel K with MRRF and the three-dimensional F_N method can readily be extended to such nonlinear inverse problems, which is an interesting future problem.

Acknowledgments

The author acknowledges support from Grant-in-Aid for Scientific Research [17K05572 and 17H02081] of the Japan Society for the Promotion of Science (JSPS) and from the JSPS A3 foresight program: Modeling and Computation of Applied Inverse Problems. This work was also supported by Hamamatsu University School of Medicine [HUSM Grant-in-Aid].

Appendix A. The F_N method in the slab geometry

We consider a slab of width z_{\max} . The radiative transport equation is given by

$$\begin{cases} \hat{\mathbf{s}} \cdot \nabla I(\mathbf{r}, \hat{\mathbf{s}}) + I(\mathbf{r}, \hat{\mathbf{s}}) = \varpi \int_{\mathbb{S}^2} p(\hat{\mathbf{s}}, \hat{\mathbf{s}}') I(\mathbf{r}, \hat{\mathbf{s}}') d\hat{\mathbf{s}}', \\ (\mathbf{r}, \hat{\mathbf{s}}) \in \mathbb{R}^3 \times \mathbb{S}^2, \\ I(\mathbf{r}, \hat{\mathbf{s}}) = f_1(\boldsymbol{\rho}, \hat{\mathbf{s}}), \quad z = 0, \quad \boldsymbol{\rho} \in \mathbb{R}^2, \quad \hat{\mathbf{s}} \in \mathbb{S}_+^2, \\ I(\mathbf{r}, \hat{\mathbf{s}}) = f_2(\boldsymbol{\rho}, \hat{\mathbf{s}}), \quad z = z_{\max}, \quad \boldsymbol{\rho} \in \mathbb{R}^2, \quad \hat{\mathbf{s}} \in \mathbb{S}_-^2, \end{cases} \quad (\text{A.1})$$

with some boundary values $f_1(\boldsymbol{\rho}, \hat{\mathbf{s}})$, $f_2(\boldsymbol{\rho}, \hat{\mathbf{s}})$. The specific intensity is given by

$$\begin{aligned} I(\mathbf{r}, \hat{\mathbf{s}}) &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q} \cdot \boldsymbol{\rho}} \sum_{m=-L}^L \left[\sum_{j=0}^{M^m-1} A^m(\nu_j^m) \right. \\ &\quad \times \mathcal{R}_{\hat{\mathbf{k}}(\nu_j^m, \mathbf{q})} \Phi_j^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu_j^m q)z/\nu_j^m} \\ &\quad + \sum_{j=0}^{M^m-1} A^m(-\nu_j^m) \mathcal{R}_{\hat{\mathbf{k}}(-\nu_j^m, \mathbf{q})} \Phi_{-j}^m(\hat{\mathbf{s}}) e^{\hat{k}_z(\nu_j^m q)z/\nu_j^m} \\ &\quad \left. + \int_{-1}^1 A^m(\nu) \mathcal{R}_{\hat{\mathbf{k}}(\nu, \mathbf{q})} \Phi_\nu^m(\hat{\mathbf{s}}) e^{-\hat{k}_z(\nu q)z/\nu} d\nu \right], \end{aligned}$$

with some coefficients $A^m(\pm\nu_j^m)$, $A^m(\nu)$. Using the orthogonality relations we have

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, 0, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(-\xi) A^m(-\xi), \end{aligned} \quad (\text{A.2})$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(-\xi) A^m(-\xi) e^{\hat{k}_z(\xi q)z_{\max}/\xi}, \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, 0, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(\xi) A^m(\xi), \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(\xi) A^m(\xi) e^{-\hat{k}_z(\xi q)z_{\max}/\xi}, \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(-\xi) A^m(-\xi) e^{\hat{k}_z(\xi q)z/\xi}, \end{aligned} \quad (\text{A.6})$$

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) I(\boldsymbol{\rho}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = 2\pi \hat{k}_z(\xi q) \mathcal{N}(\xi) A^m(\xi) e^{-\hat{k}_z(\xi q)z/\xi}. \end{aligned} \quad (\text{A.7})$$

We express $\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}})$ ($\hat{\mathbf{s}} \in \mathbb{S}^2$) as

$$\begin{aligned} \tilde{I}(\mathbf{q}, 0, -\hat{\mathbf{s}}) &= \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} c_{lm}(\mathbf{q}) Y_{lm}(\hat{\mathbf{s}}), \\ \tilde{I}(\mathbf{q}, z_{\max}, \hat{\mathbf{s}}) &= \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} d_{lm}(\mathbf{q}) Y_{lm}(\hat{\mathbf{s}}), \\ \tilde{I}(\mathbf{q}, z, -\hat{\mathbf{s}}) &= \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} b_{lm}(\mathbf{q}, z) Y_{lm}(\hat{\mathbf{s}}), \end{aligned}$$

$$\tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) = \sum_{m=-N}^N \sum_{\substack{\alpha=0 \\ l=|m|+2\alpha}}^{\lfloor (N-|m|)/2 \rfloor} a_{lm}(\mathbf{q}, z) Y_{lm}(\hat{\mathbf{s}}),$$

where $\hat{\mathbf{s}} \in \mathbb{S}_+^2$. From (A.2) and (A.3) we obtain

$$\begin{aligned} -(-1)^m \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, 0, -\hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ + \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{f}_1(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ - (-1)^m e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{f}_2(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}, \end{aligned}$$

where we used $\Phi_{-\xi}^m(\hat{\mathbf{s}}) = (-1)^m \Phi_\xi^m(-\hat{\mathbf{s}})$. Similarly from (A.4) and (A.5) we obtain

$$\begin{aligned} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ - (-1)^m \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{f}_2(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{f}_1(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ - (-1)^m e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, 0, -\hat{\mathbf{s}}) d\hat{\mathbf{s}}. \end{aligned}$$

They are rewritten as

$$\begin{aligned} \sum_{lm'} \mathcal{B}_{lm'}^m(\xi) c_{lm'} + e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \sum_{lm'} \mathcal{A}_{lm'}^m(-\xi) d_{lm'} \\ = E_{12}^m, \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \sum_{lm'} \mathcal{A}_{lm'}^m(-\xi) c_{lm'} + \sum_{lm'} \mathcal{B}_{lm'}^m(\xi) d_{lm'} \\ = E_{21}^m, \end{aligned} \quad (\text{A.9})$$

where

$$\begin{aligned} \mathcal{A}_{lm'}^m(-\xi) &= \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) Y_{lm'}(\hat{\mathbf{s}}) d\hat{\mathbf{s}}, \\ \mathcal{B}_{lm'}^m(\xi) &= (-1)^m \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) Y_{lm'}(\hat{\mathbf{s}}) d\hat{\mathbf{s}}, \\ E_{12}^m &= \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{f}_1(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &\quad + (-1)^m e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{f}_2(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}, \\ E_{21}^m &= \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{f}_2(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ &\quad + (-1)^m e^{-\hat{k}_z(\xi q)z_{\max}/\xi} \int_{\mathbb{S}_+^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{f}_1(\mathbf{q}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}. \end{aligned}$$

We obtain by using (A.2) and (A.6)

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, 0, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = e^{-\hat{k}_z(\xi q)z/\xi} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}}, \end{aligned}$$

from (A.5) and (A.7)

$$\begin{aligned} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}} \\ = e^{-\hat{k}_z(\xi q)(z_{\max}-z)/\xi} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_\xi^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}}, \end{aligned}$$

by using (A.3) and (A.6)

$$\int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} = e^{-\hat{k}_z(\xi q)(z_{\max}-z)/\xi} \times \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{-\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z_{\max}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}, \quad (\text{A.10})$$

and from (A.4) and (A.7)

$$\int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, z, \hat{\mathbf{s}}) d\hat{\mathbf{s}} = e^{-\hat{k}_z(\xi q)z/\xi} \int_{\mathbb{S}^2} \mu(\mathcal{R}_{\hat{\mathbf{k}}} \Phi_{\xi}^{m*}(\hat{\mathbf{s}})) \tilde{I}(\mathbf{q}, 0, \hat{\mathbf{s}}) d\hat{\mathbf{s}}. \quad (\text{A.11})$$

We obtain c_{lm} , d_{lm} from the linear system (A.8), (A.9). Then a_{lm} , b_{lm} are computed using (A.10), (A.11). We note that (A.10) corresponds to (5.5) and (A.11) corresponds to (5.8).

Appendix B. Forward problem

We assume point absorbers given by

$$\eta(\mathbf{r}) = \eta_a V_a \sum_{i=1}^{N_a} \delta(\mathbf{r} - \mathbf{r}_a^{(i)}),$$

where V_a corresponds to the volume of each absorber and η_a is a positive constant. We will calculate the data function using diffusion approximation. As is mentioned in (1.1), the unit of length is $\ell_t = 1/\mu_t$. With diffusion approximation we can express the Green's function as [59]

$$G(\mathbf{r}, \hat{\mathbf{s}}; \mathbf{r}', \hat{\mathbf{s}}') \simeq \frac{c\mu'_s \ell^*}{4\pi} (1 + \ell^* \hat{\mathbf{s}} \cdot \nabla)(1 - \ell^* \hat{\mathbf{s}}' \cdot \nabla') G^{(\text{DE})}(\mathbf{r}', \mathbf{r}),$$

where $\mu'_s = \varpi(1-g)$ and $\ell^* = 1/(1-\varpi g)$. Here,

$$\begin{cases} -D_0 \nabla^2 G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') + \alpha_0 G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') = \delta(\mathbf{r} - \mathbf{r}'), & z > 0, \\ G - \ell \hat{\mathbf{z}} \cdot \nabla G = 0, & z = 0, \end{cases}$$

where

$$D_0 = \frac{c\ell^*}{3}, \quad \alpha_0 = c\bar{\mu}_a,$$

with c the speed of light in the medium. We choose the extrapolation distance ($0 \leq \ell < \infty$) as

$$\ell = \frac{2}{3} \ell^*.$$

In the Fourier space we can introduce $\tilde{G}^{(\text{DE})}(z, z'; \mathbf{q}) = \tilde{G}^{(\text{DE})}(z, z'; q)$ as

$$\begin{aligned} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} e^{i\mathbf{q} \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')} \tilde{G}^{(\text{DE})}(z, z'; q) d\mathbf{q} \\ &= \frac{1}{2\pi} \int_0^\infty q J_0(q|\boldsymbol{\rho} - \boldsymbol{\rho}'|) \tilde{G}^{(\text{DE})}(z, z'; q) dq, \end{aligned} \quad (\text{B.1})$$

where J_0 is the Bessel function of the first kind of order 0. We further approximate the Green's function as

$$\tilde{G}(z, \hat{\mathbf{s}}; z', \hat{\mathbf{s}}'; \mathbf{q}) \simeq \frac{c\mu'_s \ell^*}{4\pi} (1 - i\ell^* \hat{\mathbf{s}} \cdot \mathbf{q})(1 - i\ell^* \hat{\mathbf{s}}' \cdot \mathbf{q}) \times \tilde{G}^{(\text{DE})}(z, z'; \mathbf{q}).$$

We have [60]

$$\begin{aligned} \tilde{G}^{(\text{DE})}(z, z'; q) &= \frac{1}{2D_0 Q(q)} \\ &\times \left[e^{-Q(q)|z-z'|} - \frac{1-Q(q)\ell}{1+Q(q)\ell} e^{-Q(q)|z+z'|} \right], \end{aligned}$$

where

$$Q(q) = \sqrt{k_0^2 + q^2},$$

$$k_0 = \sqrt{\frac{\alpha_0}{D_0}} = \sqrt{\frac{3\bar{\mu}_a}{\ell^*}} = \sqrt{3(1-\varpi)(1-\varpi g)}.$$

We have $k_0 \approx 0.12 \text{ mm}^{-1}$ for optical parameters in (7.1). In particular,

$$\tilde{G}^{(\text{DE})}(z, 0; q) = \frac{\ell}{D_0(1+Q(q)\ell)} e^{-Q(q)z}, \quad z > 0.$$

The integral in (B.1) can be numerically evaluated by the double-exponential formula [61]. Let $0 < j_1 < j_2 < \dots$ be zeros of J_0 . We define

$$F(q) = \tilde{G}^{(\text{DE})}(z, z'; q)q.$$

Then we have

$$\begin{aligned} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') &= \frac{1}{2\pi|\boldsymbol{\rho} - \boldsymbol{\rho}'|} \int_0^\infty J_0(x) F\left(\frac{x}{|\boldsymbol{\rho} - \boldsymbol{\rho}'|}\right) dx \\ &\approx \frac{1}{\pi|\boldsymbol{\rho} - \boldsymbol{\rho}'|} \sum_{k=1}^{N_k} J_0(q_k|\boldsymbol{\rho} - \boldsymbol{\rho}'|) F(q_k) \frac{d\phi/dy|_{y=h_j k/\pi}}{j_k (J_1(j_k))^2}, \end{aligned}$$

where $h > 0$ is a small number, N_k is an integer, $\phi(y) = y \tanh(\frac{\pi}{2} \sinh y)$, and

$$q_k = \frac{\pi}{h|\boldsymbol{\rho} - \boldsymbol{\rho}'|} \phi\left(\frac{h}{\pi} j_k\right).$$

Using the radiative transport equation, the energy density $u^{(0)}(\mathbf{r})$ is obtained as

$$u^{(0)}(\mathbf{r}) = \frac{1}{c} \int_{\mathbb{S}^2} I^{(0)}(\mathbf{r}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}.$$

Its Fourier transform has the form

$$\begin{aligned} \tilde{u}^{(0)}(\mathbf{q}, z) &= \frac{1}{c} \int_{\mathbb{S}^2} \int_{\mathbb{S}^2} \int_0^\infty \tilde{G}(z, \hat{\mathbf{s}}; z', \hat{\mathbf{s}}'; \mathbf{q}) \mu' \\ &\times \tilde{f}(\mathbf{q}, \hat{\mathbf{s}}') \delta(z') dz' d\hat{\mathbf{s}}' d\hat{\mathbf{s}}, \end{aligned}$$

where we used $\hat{\mathbf{s}}_0 = \hat{\mathbf{z}}$ and (6.1). By the diffusion approximation we have

$$\tilde{u}^{(0)}(\mathbf{q}, z) \simeq (2\pi)^2 I_0 \mu'_s \ell^* \delta(\mathbf{q} - \mathbf{q}_0) \tilde{G}^{(\text{DE})}(z, 0; \mathbf{q}_0).$$

The Fourier transform of the following $u^{(0)}$ is given by the right-hand side of the above equation.

$$u^{(0)}(\mathbf{r}) = \int_{\mathbb{R}^3} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') S(\mathbf{r}') d\mathbf{r}',$$

with the source term

$$S(\mathbf{r}) = I_0 \mu'_s \ell^* e^{i\mathbf{q}_0 \cdot \boldsymbol{\rho}} \delta(z).$$

In the diffusion approximation, $u(\mathbf{r}) = (1/c) \int_{\mathbb{S}^2} I(\mathbf{r}, \hat{\mathbf{s}}) d\hat{\mathbf{s}}$, where $I(\mathbf{r}, \hat{\mathbf{s}})$ is the solution to (1.2), satisfies

$$\begin{cases} -D_0 \nabla^2 u(\mathbf{r}) + (\alpha_0 + c\eta(\mathbf{r}))u(\mathbf{r}) = S(\mathbf{r}), & z > 0, \\ u - \ell \hat{\mathbf{z}} \cdot \nabla u = 0, & z = 0. \end{cases}$$

We have

$$\begin{aligned} u(\mathbf{r}) &= \int_{\mathbb{R}^3} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') [S(\mathbf{r}') - c\eta(\mathbf{r}')u(\mathbf{r}')] d\mathbf{r}' \\ &= u^{(0)}(\mathbf{r}) - c \int_{\mathbb{R}^3} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}') \eta(\mathbf{r}') u(\mathbf{r}') d\mathbf{r}' \\ &= u^{(0)}(\mathbf{r}) - \gamma_0 \sum_{i=1}^{N_a} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}_a^{(i)}) u(\mathbf{r}_a^{(i)}), \end{aligned} \quad (\text{B.2})$$

where

$$\gamma_0 = c\eta_a V_a.$$

Let us set

$$\eta_a = 0.0015, \quad V_a = 10^{-6} \text{ mm}^3 \mu_t^3.$$

We note that $((1/V_a) \int_{V_a} \mu_a(\mathbf{r}) d\mathbf{r})/\bar{\mu}_a \approx 4$ for each absorber. When $\mathbf{r} = \mathbf{r}_a^{(i)}$, we can write (B.2) as

$$\sum_{j=1}^{N_a} M_{ij} u(\mathbf{r}_a^{(j)}) = u^{(0)}(\mathbf{r}_a^{(i)}),$$

where

$$M_{ij} = \delta_{ij} + \gamma_0 G^{(\text{DE})}(\mathbf{r}_a^{(i)}, \mathbf{r}_a^{(j)}).$$

Thus we have

$$u(\mathbf{r}) = u^{(0)}(\mathbf{r}) - \sum_{i,j=1}^{N_a} G^{(\text{DE})}(\mathbf{r}, \mathbf{r}_a^{(i)}) T_{ij} u^{(0)}(\mathbf{r}_a^{(j)}),$$

where $T_{ij} = \gamma_0 M_{ij}^{-1}$. If $N_a = 1$, we have

$$u(\mathbf{r}) = u^{(0)}(\mathbf{r}) - \gamma_r G^{(\text{DE})}(\mathbf{r}, \mathbf{r}_a^{(1)}) u^{(0)}(\mathbf{r}_a^{(1)}),$$

where we write [62]

$$\gamma_r = T_{11} = \frac{\gamma_0}{1 + \gamma_0 G^{(\text{DE})}(\mathbf{r}_a^{(1)}, \mathbf{r}_a^{(1)})}.$$

Let us consider $G^{(\text{DE})}(\mathbf{r}_a, \mathbf{r}_a)$, where $\mathbf{r}_a = \mathbf{r}_a^{(i)}$ ($i = 1, \dots, N_a$). Introducing the ultraviolet cutoff Λ_c ($k_0 \Lambda_c \ll 1$), we define the Green's function $G^{(\text{DE})}(\mathbf{r}_a, \mathbf{r}_a)$ as [62]

$$\begin{aligned} G^{(\text{DE})}(\mathbf{r}_a, \mathbf{r}_a) &= \frac{1}{4\pi D_0} \left[\int_0^{2\pi/\Lambda_c} \frac{q}{Q(q)} dq \right. \\ &\quad \left. + \int_0^\infty \frac{q}{Q(q)} \frac{Q(q)\ell - 1}{Q(q)\ell + 1} e^{-2Q(q)z_a} dq \right] \\ &= \frac{1}{4\pi D_0} \left[Q\left(\frac{2\pi}{\Lambda_c}\right) - k_0 + \frac{e^{-2k_0 z_a}}{2z_a} \right. \\ &\quad \left. - \frac{2}{\ell} e^{2z_a/\ell} E_1\left(2z_a\left(k_0 + \frac{1}{\ell}\right)\right) \right], \end{aligned}$$

where E_1 is the exponential integral defined by

$$E_1(z) = \int_z^\infty \frac{e^{-t}}{t} dt.$$

In the numerical calculation we set

$$\Lambda_c = V_a^{1/3}.$$

The hemispheric flux is obtained as

$$\begin{aligned} J_+^{(\text{DE})}(\boldsymbol{\rho}) &= \frac{-1}{4\pi} \int_0^{2\pi} \int_{-1}^0 \mu [cu(\boldsymbol{\rho}, 0) - 3D_0 \hat{\mathbf{s}} \cdot \nabla u(\boldsymbol{\rho}, 0)] d\mu d\varphi \\ &= \frac{c}{2} \left(\frac{1}{2} + \frac{D_0}{c\ell} \right) u(\boldsymbol{\rho}, 0) = \frac{c}{2} u(\boldsymbol{\rho}, 0). \end{aligned}$$

We then have

$$\begin{aligned} J_+^{(\text{DE})(0)}(\boldsymbol{\rho}) - J_+^{(\text{DE})}(\boldsymbol{\rho}) &= \frac{c}{2} (u^{(0)}(\boldsymbol{\rho}, 0) - u(\boldsymbol{\rho}, 0)) \\ &= \frac{c}{2} \sum_{i,j=1}^{N_a} G^{(\text{DE})}(\boldsymbol{\rho}, 0, \mathbf{r}_a^{(i)}) T_{ij} u^{(0)}(\mathbf{r}_a^{(j)}). \end{aligned}$$

The data function is thus calculated as

$$\mathcal{D}(\mathbf{q}_0, \mathbf{q}) = \frac{h_d^2}{I_0} \sum_{\boldsymbol{\rho}} e^{-i(\mathbf{q}+\mathbf{q}_0)\cdot\boldsymbol{\rho}} [J_+^{(\text{DE})(0)}(\boldsymbol{\rho}) - J_+^{(\text{DE})}(\boldsymbol{\rho})]$$

$$\begin{aligned} &= \frac{c\mu_s'\ell^*}{2} \sum_{i,j=1}^{N_a} T_{ij} e^{-i\mathbf{q}\cdot\boldsymbol{\rho}_a^{(i)}} e^{-i\mathbf{q}_0\cdot(\boldsymbol{\rho}_a^{(i)} - \boldsymbol{\rho}_a^{(j)})} \\ &\quad \times \tilde{G}^{(\text{DE})}(0, z_a^{(i)}; |\mathbf{q} + \mathbf{q}_0|) \tilde{G}^{(\text{DE})}(z_a^{(j)}, 0; q_0). \end{aligned}$$

When $N_a = 1$, we have

$$\begin{aligned} \mathcal{D}(\mathbf{q}_0, \mathbf{q}) &= \frac{18\pi\eta_a V_a \mu_s' \ell^2}{4\pi\ell^* + 3\eta_a V_a c_0} e^{-i\mathbf{q}\cdot\boldsymbol{\rho}_a} \frac{e^{-Q(q_0)z_a}}{1 + Q(q_0)\ell} \\ &\quad \times \frac{e^{-Q(|\mathbf{q}+\mathbf{q}_0|)z_a}}{1 + Q(|\mathbf{q} + \mathbf{q}_0|)\ell}, \end{aligned}$$

where

$$\begin{aligned} c_0 &= Q\left(\frac{2\pi}{\Lambda_c}\right) - k_0 + \frac{e^{-2k_0 z_a}}{2z_a} \\ &\quad - \frac{2}{\ell} e^{2z_a/\ell} E_1\left(2z_a\left(k_0 + \frac{1}{\ell}\right)\right). \end{aligned}$$

We note that $c_0 \simeq Q\left(\frac{2\pi}{\Lambda_c}\right) - k_0$ for large z_a .

Supplementary material

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.jqsrt.2019.06.009.

References

- [1] Arridge SR. Optical tomography in medical imaging. *Inverse Probl* 1999;15:R41–93. doi:10.1088/0266-5611/15/2/022.
- [2] Arridge SR, Hebden JC. Optical imaging in medicine: ii. modelling and reconstruction. *Phys Med Biol* 1997;42:841–53. doi:10.1088/0031-9155/42/5/008.
- [3] Arridge SR, Schotland JC. Optical tomography: forward and inverse problems. *Inverse Probl* 2009;25:123010. doi:10.1088/0266-5611/25/12/123010.
- [4] Gibson AP, Hebden JC, Arridge SR. Recent advances in diffuse optical imaging. *Phys Med Biol* 2005;50:R1–R43. doi:10.1016/S0020-7225(98)00051-2.
- [5] Hebden JC, Arridge SR, Delpy DT. Optical imaging in medicine: i. experimental techniques. *Phys Med Biol* 1997;42:825–40. doi:10.1088/0031-9155/42/5/007.
- [6] Wang Z-M, Panasyuk GY, Markel VA, Schotland JC. Experimental demonstration of an analytic method for image reconstruction in optical diffusion tomography with large data sets. *Opt Lett* 2005;30:3338–40. doi:10.1364/OL.30.003338.
- [7] Case KM, Zweifel PF. *Linear transport theory*. Addison-Wesley; 1967.
- [8] Chandrasekhar S. *Radiative transfer*. Dover; 1960.
- [9] Cuccia DJ, Bevilacqua F, Durkin AJ, Tromberg BJ. Modulated imaging: quantitative analysis and tomography of turbid media in the spatial-frequency domain. *Opt Lett* 2005;30:1354–6. doi:10.1364/OL.30.001354.
- [10] Bassi A, D'Andrea C, Valentini G, Cubeddu R, Arridge S. Detection of inhomogeneities in diffusive media using spatially modulated light. *Opt Lett* 2009;34:2156–8. doi:10.1364/OL.34.002156.
- [11] D'Andrea C, Ducros N, Bassi A, Arridge S, Valentini G. Fast 3d optical reconstruction in turbid media using spatially modulated light. *Biomed Opt Exp* 2010;1:471–81. doi:10.1364/BOE.1.000471.
- [12] Konecky SD, Rice T, Durkin AJ, Tromberg BJ. Imaging scattering orientation with spatial frequency domain imaging. *J Biomed Opt* 2011;16:126001. doi:10.1117/1.3657823.
- [13] Zhou X, Fan Y, Hou Q, Zhao H, Gao F. Spatial-frequency-compression scheme for diffuse optical tomography with dense sampling dataset. *Appl Opt* 2013;52:1779–92. doi:10.1364/AO.52.001779.
- [14] Kaper HG. Elementary solutions of the reduced three-dimensional transport equation. *J Math Phys* 1969;10:286–97. doi:10.1063/1.1664844.
- [15] Williams MMR. The three-dimensional transport equation with applications to energy deposition and reflection. *J Phys A Math Gen* 1982;15:965–83. doi:10.1088/0305-4470/15/3/032.
- [16] Barichello LB, Siewert CE. The searchlight problem for radiative transfer in a finite slab. *J Comp Phys* 2000;157:707–26. doi:10.1006/jcph.1999.6396.
- [17] Dunn WL, Siewert CE. The searchlight problem in radiation transport: some analytical and computational results. *Z Ang Math Phys* 1985;36:581–95. doi:10.1007/BF00945298.
- [18] Kim AD. Transport theory for light propagation in biological tissue. *J Opt Soc Am A* 2004;21:820–7. doi:10.1364/JOSAA.21.000820.
- [19] Kim AD, Keller JB. Light propagation in biological tissue. *J Opt Soc Am A* 2003;20:92–8. doi:10.1364/JOSAA.20.000092.
- [20] Kim AD, Moscoso M. Beam propagation in sharply peaked forward scattering media. *J Opt Soc Am A* 2004;21:797–803. doi:10.1364/JOSAA.21.000797.
- [21] Machida M. An F_N method for the radiative transport equation in three dimensions. *J Phys A Math Theor* 2015;48:325001. doi:10.1088/1751-8113/48/32/325001.
- [22] Markel VA. Modified spherical harmonics method for solving the radiative transport equation. *Waves Random Media* 2004;14:L13–19. doi:10.1088/0959-7174/14/1/L02.

- [23] Panasyuk G, Schotland JC, Markel VA. Radiative transport equation in rotated reference frames. *J Phys A Math Gen* 2006;39:115–37. doi:10.1088/0305-4470/39/1/009.
- [24] Siewert CE. On the singular components of the solution to the searchlight problem in radiative transfer. *J Quant Spec Rad Trans* 1985;33:551–4. doi:10.1016/0022-4073(85)90022-6.
- [25] Siewert CE, Dunn WL. Radiation transport in plane-parallel media with non-uniform surface illumination. *Z Ang Math Phys* 1983;34:627–41. doi:10.1007/BF00948806.
- [26] Siewert CE, Dunn WL. The searchlight problem in radiative transfer. *J Quant Spec Rad Trans* 1989;41:467–81. doi:10.1016/0022-4073(89)90117-9.
- [27] Williams MMR. Three-dimensional transport theory: an analytical solution of an internal beam searchlight problem-i. *Ann Nucl Energy* 2009;36:767–83. doi:10.1016/j.anucene.2009.02.001.
- [28] Williams MMR. Three-dimensional transport theory: an analytical solution of an internal beam searchlight problem, iii. *Ann Nucl Energy* 2009;36:1256–61. doi:10.1016/j.anucene.2009.03.020.
- [29] Ganapol BD, Kornreich DE, Dahl JA, Nigg DW, Jahshan SN, Wemple CA. The searchlight problem for neutron in a semi-infinite medium. *Nucl Sci Eng* 1994;118:38–53.
- [30] Ganapol BD, Kornreich DE. A 3-d neutron transport benchmark solution. *Trans Theor Stat Phys* 1995;24:89–111. doi:10.1080/00411459508205121.
- [31] Ganapol BD, Kornreich DE. Three-dimensional transport theory: an analytical solution for the internal beam searchlight problem, ii. *Ann Nucl Energy* 2009;36:1242–55. doi:10.1016/j.anucene.2009.01.021.
- [32] Williams MMR. The searchlight problem in radiative transfer with internal reflection. *J Phys A Math Theor* 2007;40:6407–25. doi:10.1088/0305-4470/15/3/032.
- [33] Grandjean P, Siewert CE. The F_N method in neutron-transport theory. part ii: applications and numerical results. *Nucl Sci Eng* 1979;69:161–8. doi:10.13182/NSE79-A20608.
- [34] Siewert CE. The F_N method for solving radiative-transfer problems in plane geometry. *Astrophys Space Sci* 1978;58:131–7. doi:10.1007/BF00645380.
- [35] Benoist P, Siewert CE. The f_h method in neutron-transport theory. part i: theory and applications. *Nucl Sci Eng* 1979;69:156–60. doi:10.13182/NSE79-1.
- [36] González-Rodríguez P, Kim AD, Moscoso M. Reconstructing a thin absorbing obstacle in a half-space of tissue. *J Opt Soc Am A* 2007;24:3456–66. doi:10.1364/JOSAA.24.003456.
- [37] González-Rodríguez P, Kim AD. Reflectance optical tomography in epithelial tissues. *Inverse Probl* 2009;25:015001. doi:10.1364/JOSAA.24.003456.
- [38] Kim AD, Moscoso M. Radiative transport theory for optical molecular imaging. *Inverse Probl* 2006;22:23–42. doi:10.1088/0266-5611/22/1/002.
- [39] Liemert A, Kienle A. Light transport in three-dimensional semi-infinite scattering media. *J Opt Soc Am A* 2012;29:1475–81. doi:10.1364/JOSAA.29.001475.
- [40] Liemert A, Kienle A. Spatially modulated light source obliquely incident on a semi-infinite scattering medium. *Opt Lett* 2012;37:4158–60. doi:10.1364/OL.37.004158.
- [41] Liemert A, Kienle A. Exact and efficient solution of the radiative transport equation for the semi-infinite medium. *Sci Rep* 2013;3:1–7. doi:10.1038/srep02018.
- [42] Liemert A, Kienle A. The line source problem in anisotropic neutron transport with internal reflection. *Ann Nucl Energy* 2013;60:206–9. doi:10.1016/j.anucene.2013.05.007.
- [43] Liemert A, Kienle A. Explicit solutions of the radiative transport equation in the p_3 approximation. *Med Phys* 2014;41:111916. doi:10.1118/1.4898097.
- [44] Machida M, Panasyuk G, Schotland JC, Markel VA. The green's function for the radiative transport equation in the slab geometry. *J Phys A Math Theor* 2010;43:065402. doi:10.1088/1751-8113/43/6/065402.
- [45] Schotland JC, Markel VA. Fourier-laplace structure of the inverse scattering problem for the radiative transport equation. *Inv Prob Imag* 2007;1:181–188.
- [46] Machida M, Panasyuk GY, Wang Z-M, Markel VA, Schotland JC. Radiative transport and optical tomography with large datasets. *J Opt Soc Am A* 2016;33:551–8. doi:10.1364/JOSAA.33.000551.
- [47] Henyey LG, Greenstein JL. Diffuse radiation in the galaxy. *Astrophys J* 1941;93:70–83. doi:10.1086/144246.
- [48] V Lukic VAM, Schotland JC. Optical tomography with structured illumination. *Opt Lett* 2009;34:983–5. doi:10.1364/OL.34.000983.
- [49] Konecky SD, Mazhar A, Cuccia D, Durkin AJ, Schotland JC, Tromberg BJ. Quantitative optical tomography of sub-surface heterogeneities using spatially modulated structured light. *Opt Exp* 2009;17:14780–90. doi:10.1364/OE.17.014780.
- [50] Machida M. Singular eigenfunctions for the three-dimensional radiative transport equation. *J Opt Soc Am A* 2014;31:67–74. doi:10.1364/JOSAA.31.000067.
- [51] Garcia RDM, Siewert CE. On discrete spectrum calculations in radiative transfer. *J Quant Spec Rad Trans* 1989;42:385–94. doi:10.1016/0022-4073(89)90005-8.
- [52] Garcia RDM, Siewert CE. On computing the chandrasekhar polynomials in high order and high degree. *J Quant Spec Rad Trans* 1990;43:201–5. doi:10.1016/0022-4073(90)90052-8.
- [53] Case KM. Elementary solutions of the transport equation and their applications. *Ann Phys* 1960;9:1–23. doi:10.1016/0003-4916(60)90060-9.
- [54] McCormick NJ, Kušćer I. Bi-orthogonality relations for solving half-space transport problems. *J Math Phys* 1966;7:2036–45. doi:10.1063/1.1704886.
- [55] Mika JR. Neutron transport with anisotropic scattering. *Nucl Sci Eng* 1961;11:415–27. doi:10.13182/NSE61-1.
- [56] Dede KM. An explicit solution of the one velocity multi-dimensional boltzmann-equation in P_N approximation. *Nukleonik* 1964;6:267–71.
- [57] Kobayashi K. Spherical harmonics solutions of multi-dimensional neutron transport equation by finite fourier transformation. *J Nucl Sci Tech* 1977;14:489–501.
- [58] Machida M, Schotland JC. Inverse born series for the radiative transport equation. *Inverse Probl* 2015;31:095009. doi:10.1088/0266-5611/31/9/095009.
- [59] Markel VA, Schotland JC. Symmetries, inversion formulas, and image reconstruction for optical tomography. *Phys Rev E* 2004;70:056616. doi:10.1103/PhysRevE.70.056616.
- [60] Markel VA, Schotland JC. Inverse problem in optical diffusion tomography. ii. role of boundary conditions. *J Opt Soc Am A* 2002;19:558–66. doi:10.1364/JOSAA.19.000558.
- [61] Ogata H, Sugihara M. Quadrature formulae for oscillatory infinite integrals involving the bessel functions. *Trans J Soc Ind Appl Math* 1998;8:223–56. doi:10.11540/jjsiamt.8.2.223.
- [62] Fisher AR, Schissler AJ, Schotland JC. Photoacoustic effect for multiply scattered light. *Phys Rev E* 2007;76:036604. doi:10.1103/PhysRevE.76.036604.